

# MAIZE SILAGE RESEARCH 2025



**PIONEER**  
BRAND · PRODUCTS



# CONTENTS

**Introduction..... 1**

    Long-term breeding delivers higher yields .....2

    Interpreting the hybrid comparison t-test.....3

**Hybrid performance comparisons .....4**

    P7179.....4

    P7364.....5

    P7647.....6

    P8086.....7

    P8240.....8

    P8333.....9

    P8666.....10

    P8711.....11

    P92575.....12

    P9400.....13

    P9650.....14

    P9911.....15

    P9978.....16

    P0640.....17

    P0710.....18

    P0900.....19

    P0891.....20

    P0937.....21

    P1315.....22

    P1636.....23

    P1837.....24

**Yield and economic value of chicory.....25**







# INTRODUCTION

## Welcome to the Pioneer Maize Silage Research update for 2025.

Inside, you will find comprehensive hybrid yield data to help growers make informed decisions on which hybrid to plant. Our research programme covers more than just hybrid evaluation. Each year we conduct a range of agronomic, farm system and

environmental research projects. In this publication, we've summarised a recently published paper evaluating summer pasture and chicory yields in the Waikato.





## Long-term breeding delivers higher yields

The annual rate of maize silage yield gain in New Zealand is estimated to have been over 300 kgDM/ha/year over almost 60 years (Figure 1). Crop management and genetic improvement have both made significant contributions to yield increases.

To maximise returns, silage growers should look to introduce suitable new hybrids regularly. Desired harvest timing, soil type, cultivation methods and agronomic traits such as early growth, drought tolerance, stalk and root strength, disease resistances and silage quality are all important considerations to include in the hybrid selection process.

The most reliable way to select superior hybrids is to consider trial yield data gathered over several seasons from a wide range of locations within a growing region. An individual on-farm trial result should not be used to select a hybrid because in isolation, it is not a reliable predictor of future


hybrid performance. Hybrids should be planted and harvested at the same time. Trial data should be statistically analysed to determine if there is a real yield difference between the hybrids.

This publication provides a summary of the investment made to evaluate the silage yield performance of Pioneer® and other brands of maize silage hybrids in five defined growing regions;

**1)** Northland and north Auckland **2)** Waikato **3)** Bay of Plenty, Gisborne and northern Hawke's Bay **4)** Lower North Island and Taranaki **5)** South Island.

Summarised hybrid comparison data published in this book has been collected from field trials conducted over one or more growing seasons up to and including the 2024 harvest. The most recent regional Hybrid Performance Information (HPI) can be found at [pioneer.nz](http://pioneer.nz).

Figure 1: New Zealand maize silage yield trend



Source: New Zealand Year Book (1961 to 1996) and Pioneer® brand products New Zealand Research Programme (1997 to 2021).







# Interpreting the hybrid comparison t-test

The table below presents a summary of the possible t-test outcomes.

| P value | Confidence level | Scientific designation | Level of significance   | Yield advantage | Interpretation   |
|---------|------------------|------------------------|-------------------------|-----------------|--|
| <0.001  | >99.9%           | ★★★                    | Very highly significant | YES             | <b>Hybrid superiority for yield can be claimed.</b> Can confidently plant the winning hybrid providing no key agronomic traits are limiting. Check the trait ratings for any considerations. |
| <0.01   | >99.0%           | ★★                     | Highly significant      | YES             |  |
| <0.05   | >95.0%           | ★                      | Significant             | YES             |  |
| <0.10   | >90.0%           | CA                     | Commercially acceptable | YES             | May be regarded as a commercially acceptable basis for a decision.   |
| >0.10   | <90.0%           | NS                     | Not significant         | NO              | <b>Hybrid superiority for yield cannot be claimed.</b> Ignore the yield comparison and refer primarily to important trait ratings to select between the hybrids.                             |

The more stars (★) present for the comparison, the more confident we can be that the measured average yield difference is due to an actual genetic yield difference between the two hybrids rather than just chance.

Where a result is commercially acceptable (**CA**), the P value is <0.10 indicating the result is suitable for making a hybrid decision based on yield. Key agronomic traits must always be considered.

Where a result is not significant (**NS**), we cannot conclude there is a yield difference between the hybrids. There are two principle explanations;

1. Where the yields are very similar and the comparison has been made over a large number of locations, no significance may indicate there is little measurable difference between the two hybrids or;
2. Where there appears to be a large yield difference, no significance likely indicates there are too few trial locations, or there have been inconsistent or fluctuating results. It is therefore not possible to indicate that the difference is real.

In both instances above, growers should use regionally important hybrid trait ratings to select which hybrid to plant.

In other comparisons, yield differences may appear to be relatively small but still achieve significance – this happens in cases where yield data quality is high, and the number of trial locations is large.

**An analysis of statistical significance is carried out on all Pioneer hybrid comparisons, and we take great care to base our product yield statements and recommendations on the outcome.**







## Quickest option for the coolest growing regions.

| Feature hybrid            | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|---------------------------|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>           |                   |                  |                                       |   |                                       |
| <b>P7179</b>              | P7124             | 26               | 5.75                                  | <b>-1,659</b>                                   | ★★                                    |
| <b>P7179</b>              | P7364             | 42               | 3.29                                  | <b>-2,344</b>                                   | ★★★                                   |
| <b>P7179</b>              | P7524             | 37               | 6.58                                  | <b>-1,419</b>                                   | ★★                                    |
| <b>SOUTH ISLAND</b>       |                   |                  |                                       |   |                                       |
| <b>P7179</b>              | P7124             | 12               | 4.16                                  | <b>-1,608</b>                                   | CA                                    |
| <b>P7179</b>              | P7364             | 19               | 3.15                                  | <b>-2,510</b>                                   | ★★★                                   |
| <b>P7179</b>              | P7524             | 16               | 4.68                                  | <b>-1,694</b>                                   | ★                                     |
| <b>LOWER NORTH ISLAND</b> |                   |                  |                                       |   |                                       |
| <b>P7179</b>              | P7124             | 14               | 7.12                                  | <b>-1,703</b>                                   | ★                                     |
| <b>P7179</b>              | P7364             | 23               | 3.41                                  | <b>-2,206</b>                                   | ★★★                                   |
| <b>P7179</b>              | P7524             | 21               | 8.03                                  | <b>-1,209</b>                                   | ★                                     |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>110</b> |
| Medium yield environments      | <b>120</b> |
| High yield environments        | <b>130</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.






## The new standard for yield & earliness.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| <b>P7364</b>                             | P7179             | 42               | <b>-3.29</b>                          | 2,344   | ★★★                                   |
| <b>P7364</b>                             | P7524             | 55               | 3.58                                  | 708   | ★                                     |
| <b>P7364</b>                             | P7647             | 42               | 0.61                                  | <b>-661</b>                                     | ★                                     |
| <b>SOUTH ISLAND</b>                      |                   |                  |                                       |   |                                       |
| <b>P7364</b>                             | P7179             | 19               | <b>-3.15</b>                          | 2,510   | ★★★★                                  |
| <b>P7364</b>                             | P7524             | 24               | 1.02                                  | 904   | NS                                    |
| <b>P7364</b>                             | P7647             | 19               | 0.13                                  | <b>-875</b>                                     | NS                                    |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| <b>P7364</b>                             | P7179             | 23               | <b>-3.41</b>                          | 2,206   | ★★★                                   |
| <b>P7364</b>                             | P7524             | 31               | 5.56                                  | 556   | NS                                    |
| <b>P7364</b>                             | P7647             | 23               | 1.01                                  | <b>-484</b>                                     | NS                                    |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>110</b> |
| Medium yield environments      | <b>120</b> |
| High yield environments        | <b>130</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.





## Delivers superior yields of top-quality silage.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| <b>P7647</b>                             | P7364             | 42               | <b>-0.61</b>                          | 661   | ★                                     |
| <b>P7647</b>                             | P7524             | 37               | 2.57                                  | 1,640   | ★★★                                   |
| <b>P7647</b>                             | P8000             | 32               | 3.82                                  | 378   | NS                                    |
| <b>P7647</b>                             | P8086             | 31               | 3.56                                  | <b>-1,656</b>                                   | ★★★                                   |
| <b>SOUTH ISLAND</b>                      |                   |                  |                                       |   |                                       |
| <b>P7647</b>                             | P7364             | 19               | <b>-0.13</b>                          | 875   | NS                                    |
| <b>P7647</b>                             | P7524             | 16               | 1.17                                  | 1,961   | ★                                     |
| <b>P7647</b>                             | P8000             | 15               | 3.52                                  | 316   | NS                                    |
| <b>P7647</b>                             | P8086             | 13               | 2.91                                  | <b>-1,081</b>                                   | CA                                    |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| <b>P7647</b>                             | P7364             | 23               | <b>-1.01</b>                          | 484   | NS                                    |
| <b>P7647</b>                             | P7524             | 21               | 3.64                                  | 1,395   | ★★                                    |
| <b>P7647</b>                             | P8000             | 17               | 4.09                                  | 433   | NS                                    |
| <b>P7647</b>                             | P8086             | 18               | 4.03                                  | <b>-2,070</b>                                   | ★★★                                   |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>105</b> |
| Medium yield environments      | <b>110</b> |
| High yield environments        | <b>115</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.






CRM 80

**NEW**

## Reliable early hybrid with excellent feed value.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>SOUTH ISLAND</b>                      |                   |                  |                                       |   |                                       |
| P8086                                    | P7647             | 13               | <b>-2.91</b>                          | 1,081   | CA                                    |
| P8086                                    | P8000             | 18               | 0.43                                  | 1,106   | ★                                     |
| P8086                                    | P8240             | 18               | 3.21                                  | <b>-270</b>                                     | NS                                    |
| P8086                                    | P8333             | 18               | 1.09                                  | <b>-1,370</b>                                   | ★★                                    |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| P8086                                    | P7647             | 18               | <b>-4.02</b>                          | 2,070   | ★★★★                                  |
| P8086                                    | P8000             | 20               | <b>-1.17</b>                          | 2,266   | ★★★★                                  |
| P8086                                    | P8240             | 22               | 1.22                                  | <b>-1,041</b>                                   | CA                                    |
| P8086                                    | P8333             | 21               | <b>-0.46</b>                          | <b>-436</b>                                     | NS                                    |
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| P8086                                    | P7647             | 31               | <b>-3.56</b>                          | 1,656   | ★★★★                                  |
| P8086                                    | P8000             | 38               | <b>-0.41</b>                          | 1,717   | ★★★★                                  |
| P8086                                    | P8240             | 52               | 2.04                                  | <b>-669</b>                                     | CA                                    |
| P8086                                    | P8333             | 42               | 0.44                                  | <b>-845</b>                                     | ★★                                    |

Recommended  
growing regions



Recommended established  
plant populations (000's/ha)

Challenging yield environments **105**

Medium yield environments **115**

High yield environments **125**

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.






## Bulk and energy to fill the vat.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| P8240                                    | P8086             | 52               | -2.04                                 | 653   | CA                                    |
| P8240                                    | P8333             | 94               | -1.41                                 | 213   | NS                                    |
| P8240                                    | P8666             | 115              | 0.26                                  | 104   | NS                                    |
| <b>SOUTH ISLAND</b>                      |                   |                  |                                       |   |                                       |
| P8240                                    | P8086             | 18               | -3.21                                 | 270   | NS                                    |
| P8240                                    | P8333             | 39               | -1.24                                 | -639  | CA                                    |
| P8240                                    | P8666             | 40               | 0.27                                  | -725  | ★                                     |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| P8240                                    | P8086             | 22               | -0.45                                 | 1,041   | CA                                    |
| P8240                                    | P8333             | 37               | -1.84                                 | 1,099   | ★★                                    |
| P8240                                    | P8666             | 43               | -0.16                                 | 358   | NS                                    |
| <b>WAIKATO</b>                           |                   |                  |                                       |   |                                       |
| P8240                                    | P8086             | 12               | -0.97                                 | 517   | NS                                    |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>100</b> |
| Medium yield environments      | <b>110</b> |
| High yield environments        | <b>120</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.





CRM 83



## Highly productive mid-maturity option.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| <b>P8333</b>                             | P8086             | 42               | <b>-0.44</b>                          | 845   | ★★                                    |
| <b>P8333</b>                             | P8240             | 94               | 1.41                                  | <b>-213</b>                                     | NS                                    |
| <b>P8333</b>                             | P8666             | 135              | 1.38                                  | <b>-700</b>                                     | ★★★                                   |
| <b>P8333</b>                             | P8711             | 58               | 4.33                                  | <b>-873</b>                                     | ★★★                                   |
| <b>SOUTH ISLAND</b>                      |                   |                  |                                       |   |                                       |
| <b>P8333</b>                             | P8086             | 18               | <b>-1.09</b>                          | 1,370   | ★★                                    |
| <b>P8333</b>                             | P8240             | 39               | 1.24                                  | 639   | CA                                    |
| <b>P8333</b>                             | P8666             | 48               | 1.04                                  | <b>-413</b>                                     | NS                                    |
| <b>P8333</b>                             | P8711             | 20               | 3.98                                  | 203   | NS                                    |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| <b>P8333</b>                             | P8086             | 21               | 0.26                                  | 436   | NS                                    |
| <b>P8333</b>                             | P8240             | 37               | 1.58                                  | <b>-1,099</b>                                   | ★★                                    |
| <b>P8333</b>                             | P8666             | 60               | 1.44                                  | <b>-714</b>                                     | ★                                     |
| <b>P8333</b>                             | P8711             | 21               | 3.71                                  | <b>-713</b>                                     | CA                                    |

Recommended  
growing regions



Recommended established  
plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>100</b> |
| Medium yield environments      | <b>110</b> |
| High yield environments        | <b>115</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.





## Grows well, yields very well and feeds even better.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| P8666                                    | P8240             | 115              | -0.26                                 | -104  | NS                                    |
| P8666                                    | P8333             | 135              | -1.38                                 | 700   | ★★★                                   |
| P8666                                    | P8711             | 125              | 2.47                                  | -87   | NS                                    |
| <b>WAIKATO</b>                           |                   |                  |                                       |   |                                       |
| P8666                                    | P8240             | 28               | -0.72                                 | -879  | NS                                    |
| P8666                                    | P8333             | 25               | -2.03                                 | 1,200   | ★★                                    |
| P8666                                    | P8711             | 36               | 2.80                                  | -1,347  | ★★★                                   |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| P8666                                    | P8240             | 43               | 0.00                                  | -358  | NS                                    |
| P8666                                    | P8333             | 60               | -1.44                                 | 714   | ★                                     |
| P8666                                    | P8711             | 55               | 2.24                                  | 319   | NS                                    |
| <b>SOUTH ISLAND</b>                      |                   |                  |                                       |   |                                       |
| P8666                                    | P8240             | 40               | -0.27                                 | 725   | ★                                     |
| P8666                                    | P8333             | 48               | -1.04                                 | 413   | NS                                    |
| P8666                                    | P8711             | 28               | 1.44                                  | 911   | NS                                    |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>100</b> |
| Medium yield environments      | <b>110</b> |
| High yield environments        | <b>115</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.





## Yield with superb quality for Northern Regions.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| <b>P8711</b>                             | P8333             | 58               | <b>-4.33</b>                          | 873   | ★★                                    |
| <b>P8711</b>                             | P8666             | 125              | <b>-2.47</b>                          | 87  | NS                                    |
| <b>WAIKATO</b>                           |                   |                  |                                       |   |                                       |
| <b>P8711</b>                             | P8666             | 36               | <b>-2.80</b>                          | 1,347   | ★★★                                   |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| <b>P8711</b>                             | P8333             | 21               | <b>-3.71</b>                          | 713   | CA                                    |
| <b>P8711</b>                             | P8666             | 55               | <b>-2.24</b>                          | <b>-319</b>                                     | NS                                    |
| <b>SOUTH ISLAND</b>                      |                   |                  |                                       |   |                                       |
| <b>P8711</b>                             | P8333             | 20               | <b>-3.98</b>                          | <b>-203</b>                                     | NS                                    |
| <b>P8711</b>                             | P8666             | 28               | <b>-1.44</b>                          | <b>-911</b>                                     | NS                                    |

Recommended growing regions



Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>100</b> |
| Medium yield environments      | <b>110</b> |
| High yield environments        | <b>115</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.






## Solid, balanced hybrid, with top-of-the-line foliar health.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| P92575                                   | P9127             | 52               | -2.75                                 | 154   | NS                                    |
| P92575                                   | P9400             | 93               | -1.47                                 | 664   | ★★★                                   |
| P92575                                   | P9650             | 62               | -0.75                                 | 235   | NS                                    |
| <b>WAIKATO</b>                           |                   |                  |                                       |   |                                       |
| P92575                                   | P9127             | 25               | -3.53                                 | 787   | NS                                    |
| P92575                                   | P9400             | 50               | -1.90                                 | 985   | ★★★                                   |
| P92575                                   | P9650             | 33               | -0.66                                 | 318   | NS                                    |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| P92575                                   | P9127             | 24               | -2.05                                 | -394  | NS                                    |
| P92575                                   | P9400             | 35               | -0.91                                 | 329   | NS                                    |
| P92575                                   | P9650             | 22               | -0.62                                 | 583   | CA                                    |

Recommended growing regions



Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>95</b>  |
| Medium yield environments      | <b>110</b> |
| High yield environments        | <b>120</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.






CRM 94



## Stands tall – delivers big time.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| <b>P9400</b>                             | P92575            | 93               | <b>-0.03</b>                          | <b>-664</b>                                     | ★★★                                   |
| <b>P9400</b>                             | P9650             | 59               | 1.74                                  | <b>-522</b>                                     | CA                                    |
| <b>P9400</b>                             | P9721             | 246              | 0.81                                  | <b>-50</b>                                      | NS                                    |
| <b>WAIKATO</b>                           |                   |                  |                                       |   |                                       |
| <b>P9400</b>                             | P92575            | 50               | 1.90                                  | <b>-985</b>                                     | ★★★                                   |
| <b>P9400</b>                             | P9650             | 31               | 0.98                                  | <b>-806</b>                                     | ★                                     |
| <b>P9400</b>                             | P9721             | 109              | 2.41                                  | 1   | NS                                    |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| <b>P9400</b>                             | P92575            | 35               | 0.91                                  | <b>-329</b>                                     | NS                                    |
| <b>P9400</b>                             | P9650             | 24               | 0.44                                  | <b>-347</b>                                     | NS                                    |
| <b>P9400</b>                             | P9721             | 121              | 1.94                                  | <b>-66</b>                                      | NS                                    |

Recommended  
growing regions



Recommended established  
plant populations (000's/ha)

Challenging yield environments **100**

Medium yield environments **108**

High yield environments **115**

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.





## Security with performance.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| P9650                                    | P92575            | 62               | 0.75                                  | -235  | NS                                    |
| P9650                                    | P9400             | 59               | -0.66                                 | 522   | CA                                    |
| P9650                                    | P9911             | 42               | 2.36                                  | -1,080  | ★★                                    |
| P9650                                    | P9978             | 53               | 1.87                                  | -1,127  | ★★★                                   |
| <b>WAIKATO</b>                           |                   |                  |                                       |   |                                       |
| P9650                                    | P92575            | 33               | 0.66                                  | -318  | NS                                    |
| P9650                                    | P9400             | 31               | -0.98                                 | 806   | ★                                     |
| P9650                                    | P9911             | 21               | 1.20                                  | 91  | NS                                    |
| P9650                                    | P9978             | 24               | 1.44                                  | -978  | ★                                     |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| P9650                                    | P92575            | 22               | 0.62                                  | -583  | CA                                    |
| P9650                                    | P9400             | 24               | -0.44                                 | 347   | NS                                    |
| P9650                                    | P9911             | 21               | 3.52                                  | -2,252  | ★★★                                   |
| P9650                                    | P9978             | 25               | 2.38                                  | -1,554  | ★★★                                   |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>95</b>  |
| Medium yield environments      | <b>110</b> |
| High yield environments        | <b>120</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.



CRM 99



Optimum  
**AQUAmax**

## Top yielding, drought buster.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| <b>P9911</b>                             | P0362             | 136              | 1.03                                  | 75  | NS                                    |
| <b>P9911</b>                             | P9650             | 42               | <b>-2.36</b>                          | 1,080   | ★★                                    |
| <b>P9911</b>                             | P9721             | 258              | <b>-1.84</b>                          | 1,307   | ★★★                                   |
| <b>P9911</b>                             | P9978             | 121              | <b>-1.24</b>                          | <b>-417</b>                                     | ★                                     |
| <b>WAIKATO</b>                           |                   |                  |                                       |   |                                       |
| <b>P9911</b>                             | P0362             | 73               | 1.46                                  | 67  | NS                                    |
| <b>P9911</b>                             | P9650             | 21               | <b>-1.20</b>                          | <b>-91</b>                                      | NS                                    |
| <b>P9911</b>                             | P9978             | 53               | <b>-0.85</b>                          | <b>-612</b>                                     | ★                                     |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| <b>P9911</b>                             | P0362             | 51               | 0.49                                  | 204   | NS                                    |
| <b>P9911</b>                             | P9650             | 21               | <b>-3.52</b>                          | 2,252   | ★★★                                   |
| <b>P9911</b>                             | P9978             | 58               | <b>-1.38</b>                          | <b>-99</b>                                      | NS                                    |

Recommended  
growing regions



Recommended established  
plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>100</b> |
| Medium yield environments      | <b>108</b> |
| High yield environments        | <b>115</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.






CRM 99

## Defensive. Stable. Productive.

| Feature hybrid                           | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                          |                   |                  |                                       |   |                                       |
| P9978                                    | P0362             | 79               | 2.68                                  | -159  | NS                                    |
| P9978                                    | P9650             | 53               | -1.87                                 | 1,127   | ★★★                                   |
| P9978                                    | P9721             | 53               | -0.43                                 | 1,402   | ★★★                                   |
| P9978                                    | P9911             | 121              | 1.24                                  | 417   | ★                                     |
| <b>NORTHLAND</b>                         |                   |                  |                                       |   |                                       |
| P9978                                    | P9911             | 10               | 2.52                                  | 1,233   | ★                                     |
| <b>WAIKATO</b>                           |                   |                  |                                       |   |                                       |
| P9978                                    | P0362             | 44               | 2.86                                  | -473  | CA                                    |
| P9978                                    | P9650             | 24               | -1.44                                 | 978   | ★                                     |
| P9978                                    | P9911             | 53               | 0.85                                  | 612   | ★                                     |
| <b>LOWER NORTH ISLAND &amp; TARANAKI</b> |                   |                  |                                       |   |                                       |
| P9978                                    | P9650             | 30               | 2.12                                  | 220   | NS                                    |
| P9978                                    | P9721             | 25               | -2.38                                 | 1,554   | ★★★                                   |
| P9978                                    | P9911             | 58               | 1.38                                  | 99  | NS                                    |

Recommended growing regions



Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>95</b>  |
| Medium yield environments      | <b>110</b> |
| High yield environments        | <b>120</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.



## Leaf disease champion delivering silage yield stability.

| Feature hybrid                        | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|---------------------------------------|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                       |                   |                  |                                       |   |                                       |
| <b>P0640</b>                          | P0362             | 80               | <b>-1.09</b>                          | 1,077   | ★★★                                   |
| <b>P0640</b>                          | P0900             | 85               | 0.60                                  | 662   | ★                                     |
| <b>P0640</b>                          | P0937             | 91               | 1.78                                  | 1,134   | ★★★                                   |
| <b>P0640</b>                          | P1315             | 37               | 1.58                                  | <b>-38</b>                                      | NS                                    |
| <b>NORTHLAND &amp; SOUTH AUCKLAND</b> |                   |                  |                                       |   |                                       |
| <b>P0640</b>                          | P0891             | 29               | <b>-2.76</b>                          | <b>-262</b>                                     | NS                                    |
| <b>P0640</b>                          | P0900             | 12               | <b>-0.77</b>                          | 433   | NS                                    |
| <b>P0640</b>                          | P0937             | 12               | 0.10                                  | 701   | NS                                    |
| <b>WAIKATO</b>                        |                   |                  |                                       |   |                                       |
| <b>P0640</b>                          | P0362             | 58               | <b>-0.79</b>                          | 1,417   | ★★★                                   |
| <b>P0640</b>                          | P0900             | 65               | 0.97                                  | 891   | ★★                                    |
| <b>P0640</b>                          | P0937             | 75               | 1.86                                  | 1,295   | ★★★                                   |
| <b>P0640</b>                          | P1315             | 30               | 2.01                                  | <b>-34</b>                                      | NS                                    |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>95</b>  |
| Medium yield environments      | <b>105</b> |
| High yield environments        | <b>110</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.



**NEW**

optimum  
**AQUAmax**



CRM 107

## Exceptional foliar health and yield stability – wet or dry!

| Feature hybrid  | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|-----------------|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b> |                   |                  |                                       |   |                                       |
| <b>P0710</b>    | P0900             | 48               | 1.43                                  | <b>-290</b>                                     | NS                                    |
| <b>P0710</b>    | P0937             | 35               | 2.59                                  | 457   | CA                                    |

Recommended  
growing regions



Recommended established  
plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>85</b>  |
| Medium yield environments      | <b>95</b>  |
| High yield environments        | <b>115</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.



## Reliable veteran.

| Feature hybrid   | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|------------------|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>  |                   |                  |                                       |   |                                       |
| <b>P0891</b>     | P0640             | 143              | 1.45                                  | 6   | NS                                    |
| <b>P0891</b>     | P0900             | 82               | 1.62                                  | -451  | NS                                    |
| <b>P0891</b>     | P0937             | 113              | 3.12                                  | -25   | NS                                    |
| <b>P0891</b>     | P1315             | 27               | 2.55                                  | -2,245  | ★★★                                   |
| <b>NORTHLAND</b> |                   |                  |                                       |   |                                       |
| <b>P0891</b>     | P0640             | 29               | 2.76                                  | 262   | NS                                    |
| <b>P0891</b>     | P0900             | 13               | 1.65                                  | 539   | NS                                    |
| <b>P0891</b>     | P0937             | 19               | 3.14                                  | 1,481   | ★★★                                   |
| <b>WAIKATO</b>   |                   |                  |                                       |   |                                       |
| <b>P0891</b>     | P0640             | 115              | 0.98                                  | -394  | CA                                    |
| <b>P0891</b>     | P0900             | 68               | 1.39                                  | -843  | ★                                     |
| <b>P0891</b>     | P0937             | 89               | 2.66                                  | -543  | ★                                     |
| <b>P0891</b>     | P1315             | 36               | 2.34                                  | -2,234  | ★★★                                   |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>95</b>  |
| Medium yield environments      | <b>105</b> |
| High yield environments        | <b>110</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.






## A proven, stable, all-round hybrid.

| Feature hybrid       | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|----------------------|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>      |                   |                  |                                       |   |                                       |
| P0900                | P0640             | 85               | -0.60                                 | -662  | ★                                     |
| P0900                | P0710             | 48               | -1.03                                 | -290  | NS                                    |
| P0900                | P0937             | 117              | 1.24                                  | 285   | NS                                    |
| P0900                | P1315             | 53               | 0.57                                  | -448  | NS                                    |
| P0900                | P1636             | 50               | 2.48                                  | -1,588  | ★★★                                   |
| <b>NORTHLAND</b>     |                   |                  |                                       |   |                                       |
| P0900                | P0640             | 12               | 0.69                                  | -433  | NS                                    |
| P0900                | P0937             | 17               | 1.59                                  | -389  | NS                                    |
| <b>WAIKATO</b>       |                   |                  |                                       |   |                                       |
| P0900                | P0710             | 40               | -1.30                                 | 193   | NS                                    |
| P0900                | P0891             | 68               | -1.39                                 | 843   | ★★                                    |
| P0900                | P0937             | 88               | 0.96                                  | 520   | ★                                     |
| P0900                | P1315             | 49               | 0.61                                  | -550  | CA                                    |
| <b>BAY OF PLENTY</b> |                   |                  |                                       |   |                                       |
| P0900                | P0937             | 11               | 2.85                                  | -537  | NS                                    |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>85</b>  |
| Medium yield environments      | <b>95</b>  |
| High yield environments        | <b>115</b> |


<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.



## Solid hybrid with great standability and foliar health.

| Feature hybrid                                   | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                                  |                   |                  |                                       |   |                                       |
| P0937  | P0640             | 91               | -1.78                                 | -1,134  | ★★★                                   |
| P0937  | P0710             | 35               | -2.59                                 | -457  | CA                                    |
| P0937  | P0900             | 117              | -1.24                                 | -285  | NS                                    |
| P0937  | P1315             | 57               | -0.69                                 | -1,504  | ★★★                                   |
| <b>NORTHLAND</b>                                 |                   |                  |                                       |   |                                       |
| P0937  | P0640             | 12               | -0.98                                 | -701  | NS                                    |
| P0937  | P0891             | 19               | -3.14                                 | -1,481  | ★★                                    |
| P0937  | P0900             | 17               | -1.59                                 | 389   | NS                                    |
| <b>WAIKATO</b>                                   |                   |                  |                                       |   |                                       |
| P0937  | P0640             | 75               | -1.86                                 | -1,295  | ★★★                                   |
| P0937  | P0900             | 88               | -0.96                                 | -520  | ★                                     |
| P0937  | P1315             | 51               | -0.41                                 | -1,338  | ★★★                                   |
| <b>BAY OF PLENTY, GISBORNE &amp; HAWKE'S BAY</b> |                   |                  |                                       |   |                                       |
| P0937  | P0900             | 11               | -2.85                                 | 537   | NS                                    |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>90</b>  |
| Medium yield environments      | <b>100</b> |
| High yield environments        | <b>115</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.






## Defensive from Northland to Hawke's Bay.

| Feature hybrid       | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|----------------------|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>      |                   |                  |                                       |   |                                       |
| <b>P1315</b>         | P1185             | 9                | 0.74                                  | <b>-765</b>                                     | NS                                    |
| <b>P1315</b>         | P1636             | 25               | 2.11                                  | 257   | NS                                    |
| <b>WAIKATO</b>       |                   |                  |                                       |   |                                       |
| <b>P1315</b>         | P0891             | 36               | <b>-2.34</b>                          | 2,234   | ★★★                                   |
| <b>P1315</b>         | P0900             | 49               | <b>-0.61</b>                          | 550   | CA                                    |
| <b>P1315</b>         | P0937             | <b>51</b>        | 0.41                                  | 1,338   | ★★★                                   |
| <b>P1315</b>         | P1636             | <b>58</b>        | 1.88                                  | <b>-774</b>                                     | ★                                     |
| <b>BAY OF PLENTY</b> |                   |                  |                                       |   |                                       |
| <b>P1315</b>         | P1477W            | 25               | 2.51                                  | <b>-785</b>                                     | NS                                    |
| <b>P1315</b>         | P1636             | 25               | 1.81                                  | <b>-519</b>                                     | NS                                    |

Recommended growing regions



Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>80</b>  |
| Medium yield environments      | <b>90</b>  |
| High yield environments        | <b>100</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.



## Enjoy the agronomics of this top-yielding hybrid.

| Feature hybrid                                   | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                                  |                   |                  |                                       |   |                                       |
| <b>P1636</b>                                     | P0900             | 50               | <b>-2.48</b>                          | 1,588   | ★★★                                   |
| <b>P1636</b>                                     | P0937             | 66               | <b>-1.03</b>                          | 1,979   | ★★★                                   |
| <b>P1636</b>                                     | P1315             | 108              | <b>-1.92</b>                          | 476   | CA                                    |
| <b>P1636</b>                                     | P1477W            | 155              | 1.11                                  | <b>-474</b>                                     | ★                                     |
| <b>P1636</b>                                     | P1837             | 53               | 1.62                                  | 681   | ★                                     |
| <b>NORTHLAND &amp; SOUTH AUCKLAND</b>            |                   |                  |                                       |   |                                       |
| <b>P1636</b>                                     | P1185             | 7                | <b>-1.34</b>                          | <b>-1,679</b>                                   | CA                                    |
| <b>P1636</b>                                     | P1315             | 25               | <b>-2.11</b>                          | <b>-257</b>                                     | NS                                    |
| <b>P1636</b>                                     | P1477W            | 41               | 0.11                                  | <b>-873</b>                                     | CA                                    |
| <b>P1636</b>                                     | P1837             | 13               | 1.27                                  | 859   | NS                                    |
| <b>WAIKATO</b>                                   |                   |                  |                                       |   |                                       |
| <b>P1636</b>                                     | P0900             | 46               | <b>-2.57</b>                          | 1,761   | ★★★                                   |
| <b>P1636</b>                                     | P0937             | 55               | <b>-1.32</b>                          | 2,134   | ★★★                                   |
| <b>P1636</b>                                     | P1315             | 58               | <b>-1.88</b>                          | 774   | ★                                     |
| <b>P1636</b>                                     | P1477W            | 75               | 1.80                                  | <b>-125</b>                                     | NS                                    |
| <b>P1636</b>                                     | P1837             | 30               | 2.07                                  | 683   | NS                                    |
| <b>BAY OF PLENTY, GISBORNE &amp; HAWKE'S BAY</b> |                   |                  |                                       |   |                                       |
| <b>P1636</b>                                     | P1315             | 25               | <b>-2.31</b>                          | 519   | NS                                    |
| <b>P1636</b>                                     | P1477W            | 39               | <b>-1.92</b>                          | <b>-725</b>                                     | NS                                    |
| <b>P1636</b>                                     | P1837             | 10               | 0.91                                  | 446   | NS                                    |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |            |
|--------------------------------|------------|
| Challenging yield environments | <b>95</b>  |
| Medium yield environments      | <b>105</b> |
| High yield environments        | <b>110</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.





## Defensive full-season giant.

| Feature hybrid                                   | Comparison hybrid | Number of trials | Drymatter difference (%) <sup>1</sup> | Yield advantage to P7364 (kgDM/ha) <sup>2</sup> | Statistical significance <sup>3</sup> |
|--|-------------------|------------------|---------------------------------------|---|---------------------------------------|
| <b>NATIONAL</b>                                  |                   |                  |                                       |   |                                       |
| P1837  | P1477W            | 54               | -0.22                                 | -1,226  | ★★                                    |
| P1837  | P1636             | 53               | -1.62                                 | -681  | ★                                     |
| <b>WAIKATO</b>                                   |                   |                  |                                       |   |                                       |
| P1837  | P1477W            | 33               | -0.07                                 | -987  | CA                                    |
| P1837  | P1636             | 30               | -2.07                                 | -683  | NS                                    |
| <b>BAY OF PLENTY, GISBORNE &amp; HAWKE'S BAY</b> |                   |                  |                                       |   |                                       |
| P1837  | P1477W            | 11               | -1.53                                 | -1,252  | NS                                    |
| P1837  | P1636             | 13               | -1.27                                 | -859  | NS                                    |

### Recommended growing regions



### Recommended established plant populations (000's/ha)

|                                |           |
|--------------------------------|-----------|
| Challenging yield environments | <b>70</b> |
| Medium yield environments      | <b>80</b> |
| High yield environments        | <b>90</b> |

<sup>1</sup>Positive DM differences means the bolded hybrid was drier at harvest, negative DM differences mean it was wetter. <sup>2</sup>A positive yield advantage means the bolded hybrid produced more yield, a negative yield advantage means it produced less. <sup>3</sup>For information on interpreting hybrid comparison data and statistical significance see page 3.

# YIELD AND ECONOMIC VALUE OF CHICORY GROWN ON FIVE WAIKATO DAIRY FARMS<sup>1</sup>

## Introduction

Chicory has become a popular forage crop for dairy farmers due to its high expected yield and nutritional benefits for livestock over the summer months. Outside of being part of a pasture renewal program, farmers have opted to grow chicory to supply quality feed (energy and protein) during the summer months when pasture yield and quality is expected to decline on most dairy farms.

However, there is little information available to quantify the drymatter yield of pure chicory swards vs pasture when included in a grazed farm system. Given the costs associated with establishing chicory, and that farmers ideally require a minimum of 10% of the farm area in chicory to supply >3kg DM/cow, it is important to know whether chicory is yielding as expected and improving farm productivity.

The objective of this trial was to determine chicory and pasture yields from October to May in dairy systems and to consider the cropping costs against alternative options to supply summer feed.

## Method

The study was conducted during the 2022-23 and 2023-24 seasons on five commercial dairy farms between Hamilton and Te Awamutu in the Waikato. The farms varied in scale, system intensity and proportion of chicory inclusion (Table 1). Chicory crops were established between the 28th September and 20th October and sprayed out between the 12th March and 19th April of the following year before pasture renewal. Chicory was primarily strip grazed 3-6 times between December and April, providing 1 to 3 kg DM of the daily dairy cow diet during the summer period.

| Farm | Farm area (ha) | Number of cows | Stocking rate | Chicory area (ha) | Farm system (1-5) |
|------|----------------|----------------|---------------|-------------------|-------------------|
| A    | 52             | 165            | 3.2           | 5                 | 3                 |
| B    | 124            | 415            | 3.4           | 15                | 5                 |
| C    | 213            | 850            | 4.0           | 36                | 5                 |
| D    | 140            | 535            | 3.8           | 7                 | 4                 |
| E    | 92             | 316            | 3.4           | 4                 | 2                 |

Table 1: Trial farm system details.

In spring, designated chicory paddocks were sprayed out, and five herbage cages (1 m<sup>2</sup>) were placed onto the chicory paddocks after they were sown. Five herbage cages were placed on an adjacent permanent pasture paddock that represented average pasture production on the farm. These cages prevented dairy cows from grazing, allowing for the assessment of drymatter production without animal interference. After each grazing event of the pasture paddocks, the vegetation under each cage was mown to a height of 4 cm using a standard lawnmower. Once the chicory paddocks were established and had grown sufficiently to be grazed (December-April), the same measurement method was applied.

Cut grass and chicory were collected and weighed to determine the fresh weight. A subsample was submitted to a commercial laboratory for drymatter assessment. After samples were

collected, cages were moved onto the grazed area so regrowth was representative of the grazed area.

Chicory cuts finished when the paddock was sprayed out. Pasture cage cuts were taken at every grazing event until 1st May, when the chicory paddocks returned to the grazing rotation.


## Results

During the 2022/23 summer, January-March rainfall was 369 mm (Waikato Regional Council environmental data hub), 185% of the long-term average (199mm). In the 2023/24 summer, Jan-Mar rainfall dropped back to 219 mm, 110% of the long-term average. Consequently, monthly pasture growth rates (Jan-Mar) for the 2022/23 and 2023/24 seasons were consistently higher than the expected average (based on pasture growth rate data from nearby Owl Farm).




## Growth rates

On average, the chicory area produced substantially less yield than perennial ryegrass during the establishment phase from October to December and a similar yield from January to March (Figure 1).



**Figure 1:** Monthly pasture and chicory yield, average over 2022/23 and 2023/24.

Over the two years of the trial, chicory produced an average of 6,771 kg DM/ha, while perennial ryegrass yielded an average of 11,508 kg DM/ha, meaning that chicory yielded around 4,700 kg DM/ha less than pasture per year (Figure 2).



**Figure 2:** Pasture and chicory yield results Oct-May.

# Financial

Over the two years of the trial, the average on-farm establishment cost for chicory was \$1,052/ha and ranged between \$740–\$1,582/ha (Table 2). The most notable drivers behind the wide range in

establishment costs were whether starter fertiliser was applied, the post-emergent weed control program and in the case of Farm D, having to replant the chicory crop due to failed germination.

|        | Year  | Spray out (\$/ha) | Fertiliser (\$/ha) | Planting (\$/ha) | Post-emergence weed control (\$/ha) | Total cost (\$/ha) | Chicory yield (tDM/ha) |
|--------|-------|-------------------|--------------------|------------------|-------------------------------------|--------------------|------------------------|
| Farm A | 22/23 | \$135.12          | \$243.05           | \$634.80         | \$0.00                              | \$1,012.97         | 6.2                    |
|        | 23/24 | \$115.00          | \$119.50           | \$676.85         | \$0.00                              | \$911.35           | 7.4                    |
| Farm B | 22/23 |                   |                    |                  |                                     |                    | 5.2                    |
|        | 23/24 | \$96.16           | \$130.00           | \$581.60         | \$131.00                            | \$938.76           | 6.7                    |
| Farm C | 22/23 | \$114.85          | \$0.00             | \$530.55         | \$94.79                             | \$740.19           | 4.6                    |
|        | 23/24 | \$92.15           | \$0.00             | \$655.44         | \$97.20                             | \$844.79           | 6.1                    |
| Farm D | 22/23 | \$113.23          | \$0.00             | \$475.44         | \$371.10                            | \$959.77           | 6.3                    |
|        | 23/24 | \$110.11          | \$0.00             | \$870.42         | \$601.19                            | \$1,581.72         | 8.6                    |
| Farm E | 22/23 |                   |                    |                  |                                     |                    | 8.5                    |
|        | 23/24 | \$161.50          | \$258.65           | \$859.50         | \$147.00                            | \$1,426.65         | 8.0                    |

Table 2: Breakdown of per-hectare establishment costs.





## Discussion and conclusions

Farmers reasons for growing chicory include managing spring pasture surpluses, feeding youngstock, improving summer feed quantity and quality, weed control and assisting with pasture renewal.

During the two trial seasons, growing chicory shifted a surplus of feed from spring to summer/autumn but did not increase total drymatter production above that achieved from a well-managed ryegrass-clover pasture.

Long-term average pasture analysis data collected from ten Waikato dairy farms between January and March over 12 seasons shows an average energy content of 11.2 MJ/kg DM and crude protein of 24.8% (pers comm. Tim Sandbrook, Open Country Dairy). While this is slightly less energy (12.5 MJ/kg DM) and more protein (20.6%) than previously measured in chicory, it is still sufficient to support mid-to-late lactation milk production.

When the opportunity cost of lost pasture production was considered, chicory became uneconomic because it produced less drymatter than the pasture.

When considering feed quantity, quality and establishment costs over the two years of the trial, farmers would have been better off to use the money invested in chicory to purchase a high-quality protein

concentrate (e.g., dried distiller's grain) to provide additional energy and protein if required. Imported feed is more flexible and could be used when needed, at the required rates to strategically fill feed gaps without displacing pasture production.

The improvement in pasture production as a flow-on effect from growing a chicory crop was not included in the scope of this trial. If pastures had “run out” and were more than 40% below full production, then growing chicory followed by sowing new pasture would have a more positive impact on profitability, but would need to be balanced against alternative methods of pasture renewal. To ensure the best chance of a higher-yielding chicory crop, establishment costs should not be skimped.

Based on this analysis, for chicory to be an economically viable crop, farmers would have to have experienced a 30% reduction in pasture production (Oct-May) and a 30% lift in chicory production. This is a significant yield gap to bridge.

## Acknowledgements

We greatly appreciate the five farmers involved and thank Logan Dawson, Andy Macky, Andrew McPherson, Dave Swney and David Warren for their time and feedback throughout the trial. Thanks also to David Baird for assistance with statistical analysis of the data.

<sup>1</sup>Bell, W.; Journeaux, P. 2024. Yield and economic value of chicory grown on five Waikato dairy farms. Proceedings of the Australasian Dairy Science Symposium 2024







**talk to us**  
**0800 746 633**



**more online at**  
**pioneer.nz**



**[youtube.com/pioneerbrandproductsnz](https://youtube.com/pioneerbrandproductsnz)**



**[facebook.com/pioneerbrandproducts](https://facebook.com/pioneerbrandproducts)**

**Genetic Technologies Limited, Gisborne Office**

328 Lytton Road, PO Box 214, Gisborne 4040. Phone: 06 869 0660 Fax: 06 867 0083

Pioneer® brand products are provided subject to the terms and conditions of purchasing, which are part of the labelling and purchase documents.  
®. TM. SM Trademarks and service marks of Dow AgroSciences, DuPont or Pioneer, and their affiliated companies or their respective owners. ©2025 Corteva. ©2025, Genetic Technologies Limited. All Rights Reserved.

The information in this publication is general in nature only. Although the information in this publication is believed to be accurate, no liability (whether as a result of negligence or otherwise) is accepted for any loss of any kind that may arise from actions based on the contents of this publication. ©2025, Genetic Technologies Limited. No part of this publication can be reproduced without prior written consent from Genetic Technologies Limited.



**PIONEER®**  
BRAND · PRODUCTS