PIONEER

ov Silage

A YATES FAMILY BUSINESS

2025-2026


Like father like son

Maize silage the answer for generations of Swneys

Five challenges facing dairy farm systems Taking good care of your maize seed







t is a pleasure to share with you the Pioneer[®] Maize for Silage catalogue for 2025 which contains details of our latest hybrid lineup, technical articles on looking after your maize seed, and the value of maize silage in dairy farm systems. We are grateful to the Singh, Swney and Bailey families who have allowed us to showcase their successful dairy farm businesses.

This year marks the 50th anniversary of the signing of the seed production and distribution agreement between Pioneer Hi-Bred and my father, Philip Yates, who was the Managing Director of Arthur Yates and Co. Ltd at the time. It's a great privilege for our family to have represented the Pioneer brand in the New Zealand market for the past fifty years. I want to sincerely thank all the farmers who have planted Pioneer maize seed as well as the merchants, contractors, consultants and the Pioneer team who have supported them over all those years.

We are pleased to introduce P0710 and P1185 to our lineup this season. P0710 (page 20) is our newest Optimum[®] AQUAmax[®] hybrid. These resilient products are a great option for growers planting on lighter soils or in lower rainfall areas. P1185 (page 22) is a foliar health champion with proven Northern Leaf Blight and Rust resistances.

It's been pleasing to see global dairy prices improve and interest rates drop over the past year bringing welcome relief to farmers who have been challenged by high operating and finance costs and lower product returns. Higher milk prices represent a great opportunity for farmers to further develop their farm systems to increase milk production and profit, and maize silage has an important role to play. Not only is it cost-effective, but it also offers sustainability advantages, especially when compared to imported feeds. If you are considering making changes to your farm system, feel free to contact one of our Dairy Specialists, Wade Bell or Matt Dalley, for an obligation-free visit.

The Pioneer team joins me in wishing you all the very best for the 2025-26 season. As always, we appreciate your support for Pioneer[®] brand products and are committed to helping you extract the maximum value from them. If we can help you in any way, please give us a call.

2025

With warmest regards,

within fale

William Yates Managing Director

20 20



MAIZE FOR SILAGE 2025-2026

Farm success stories

10 Maize runs in the family

COVER STORY

- 34 Maize helps power production
- 52 A proud history, an exciting future

Inside Pioneer

- 4 Feed of the future
- 24 LumiGEN® seed treatments
- 28 Looking after your maize seed
- $38 \quad \text{Pioneer celebrates 50 years in NZ} \\$

Pioneer[®] brand products

- 14 Maize hybrids
- 46 Inoculants
- 51 Summer feed

Calculators & guides

- 56 Growing and harvest cost guide
- 57 Maize silage drymatter cost
- 58 Trait characteristic notes
- 59 Choosing the right hybrid for your farm
- 60 Hybrid options for your region
- 62 Hybrid trait characteristics





7/mgfadese



PIONEER LONG LOOK

We strive to produce the best products on the market.

We deal honestly and fairly with customers, employees and business associates.

We vigorously market our products, but without misrepresentation.

We provide helpful management information to assist customers in making optimum profits from our products.





Feed of the later of the later



he future for dairying looks increasingly bright with research continuing to highlight the nutritional benefits of milk products. However, farm systems will need to adapt to a changing climate as well as increased consumer demands for improved cow welfare and better environmental stewardship.



We asked Pioneer Farm Systems Specialists Matt Dalley and Wade Bell what they see as the five key challenges for dairy farm systems now and into the future, and the role of maize silage and grain in future feed systems. Here's what they said.

Pasture harvest

In many districts pasture yields are stable or even decreasing and there are increasing challenges to pasture persistence. We can expect climate change to deliver greater variation in rainfall and seasonal pasture growth rates. An added challenge will be feed quality. Under warm, dry summer conditions, perennial ryegrass will become less competitive than less nutritious C4 grasses including paspalum and kikuyu.

Why maize?

- Feed maize silage to prevent overgrazing, creating positive substitution and increasing pasture harvest.
- Use maize silage followed by a winter crop to double the drymatter yield on a percentage of your farm and increase the yield of homegrown feed.
- Couple maize silage with a standoff facility to keep cows off wet pastures minimising pugging damage to pastures.
- Utilise maize silage as an integral part of a successful pasture renewal programme.



Reproductive performance

Empty rates and six week in-calf rates continue to be a big challenge for a number of the farms we visit. It's a big problem for the industry because it drives the need for higher replacement rates and impacts a farmer's ability to cull cows based on their genetic merit and other factors like age, health status, conformation or temperament. While the reasons for poor in-calf rates are numerous and complex, nutrition plays an important role.

Fertility is affected by body condition score (BCS) at calving, BCS at the planned start of mating, loss of BCS in early lactation, and whether cows are gaining liveweight before and during mating¹.

Research showed that NZ Holstein-Friesian cows fed 3 or 6 kgDM of a maize grain-based supplement lost less condition and started to gain condition faster than cows fed pasture only diets' (Figure 1).



Why maize?

- Maize silage is an excellent feed for reducing summer condition score loss and putting condition on cows during the autumn months.
- Maize is the highest energy grain which makes it ideal for increasing energy intake in early lactation reducing condition score loss.





Figure 1: Body condition scores change for cows fed varying rates of a maize grain-based supplement¹.

¹ Body condition scoring – the reference guide for New Zealand farmers https://www.dairynz.co.nz/media/hjpl5oj2/body-condition-score-reference-guide-2024.pdf



D3. Per cow production

New Zealand dairy cow production has historically been lower than that achieved in Australian, Irish or North American farm systems. While per cow production has slowly risen over time, the average farm produces around 400 kgMS/cow or around 0.8 kgMS per kilogram of liveweight².

Many of the farms we visit are focused on lifting per cow performance to help improve the efficiency of their system. The keys to high per cow production are a condensed calving, a high peak and a slow decline so that days in milk are maximised.

² Assumes an average cow liveweight of 500 kg

Why maize?

- In spring cows can often eat more drymatter when they are fed maize silage as well as low drymatter pasture.
- Very little energy is expended eating 3-5 kgDM on a feedpad versus grazing the equivalent amount of pasture drymatter in the paddock.
- Carrying a buffer of maize silage allows farmers to fill deficits where feed demand exceeds pasture growth.
- Maize is ideal for extending lactation while at the same time meeting cow condition and pasture cover targets.

04. Labour efficiency

Often when we ask farmers what their key challenges are, labour features somewhere near the top of the list. There is a shortage of skilled workers in the dairy industry and the challenge of finding and retaining good staff seems to increase the further you live from a major town or city. We see an increasing number of farmers who are investing in cow collar technology and cow housing and feeding infrastructure mainly because they save labour and/or make cow management simpler. Lifting per cow performance is another way to increase labour efficiency because less animals need to be milked and less replacements raised for the same milk output.

Why maize?

- Well-designed farm systems combining maize silage and a feedpad can help improve labour efficiency.
- Large amounts of maize silage can be fed out quickly and farmers often comment that feeding maize silage is easier and less risky than managing the intake of some grazed forage crops.
- The most efficient systems combine timed gate latches or wearable technology with well-designed feeding facilities located close to silage storage areas. Cows make their own way to the feedpad offering significant time savings especially early in the morning.

Environmental sustainability

Whether we like it or not, global food company demand for sustainably produced milk will drive milk pricing and ultimately on-farm practices. Chief Executive Miles Hurrell told the Farmers Weekly In Focus podcast that Fonterra's Scope 3 emissions programme is vital if the co-operative wants to continue to supply key customers³. Ninety percent of these emissions occur on farm⁴ and maize silage can help by improving animal efficiency (see Point 3 on page 7), optimising nitrogen use and reducing a farmer's reliance on imported feed.

Why maize?

- Maize has high nitrogen use efficiency a kilogram of nitrogen fertiliser can produce around 55 kgDM of maize silage but around 15 kgDM of pasture⁵.
- Growing maize silage on effluent paddocks is an excellent way to utilise surplus nitrogen whilst producing low cost drymatter.
- Maize silage has a beneficial impact on nitrous oxide emissions due to its lower nitrogen concentration than pasture⁵.
- Feeding 5 kgDM of maize silage will give you a lower Scope 3 greenhouse gas emission than feeding 1 kgDM of palm kernel extract.

Grain based 2.0 Pasture based 1.0

PKE 5.8

By-products 0.03

Figure 2: Relative greenhouse gas emissions associated with supplementary feed⁴.

³ https://www.farmersweekly.co.nz/markets/scope-3-work-is-fonterras-ticket-toride/#google_vignette ⁴ Fonterra, 2023. Our approach to on-farm emissions. May 2023

Williams et al. 2010. Using maize silage to reduce the impact of dairy farm systems on water use and quality in New Zealand: A review. Proceedings of the Australasian Dairy Science Symposium pp 74-77

Hear more from our Farm Systems Specialists

"Feed for Thought" is a regular podcast where Pioneer® Farm Systems Specialists Wade Bell

and Matt Dalley share practical and relevant information on technical and farm systems topics. Our aim is to prompt thought, discussion and actions which improve the profitability and sustainability of farming. Tune in and listen to the team as well as our guests who include farmers, researchers, agronomists and nutritionists from New Zealand and abroad. **To listen and subscribe to Feed for Thought, scan here:**



FARM SUCCESS STORIES

Maize runs in the lawing

DAVE SWNEY AND ALICE TREVELYAN, TE KAWA



or more than 30 years maize silage has underpinned the profitability and sustainability of Waikato-based Ivondale farm.

The original 43 ha farm located at Te Kawa, just south of Te Awamutu, was purchased by Phill and Deb Swney in 1990 and was slowly expanded through the lease and purchase of adjoining land.

The current 123 ha (eff) dairy operation is run by son Dave and his partner Alice Trevelyan who own 50% of the assets and contract milk for the business.

In the 2023-24 season, the 410-cow crossbred herd produced 189,500 kgMS, equating to 1,541 kgMS/ha or 462 kgMS/cow. Farm working expenses were \$5.85/kgMS and the property had an operating profit of \$4,893/ha.

This season (2024-25) the farm is on track to produce just over 200,000 kgMS from 410 cows.

Dave, who was a DairyNZ Consulting Officer for five years, counts himself lucky to have come home to a farm with good infrastructure and a well-bred Friesian x Jersey (F8J8) herd.

"Dad bred an amazing herd of big black crossbreds that combine the best of both breeds" says Dave. "We feel really lucky to be able to carry the genetics on".

Determined to increase production and believing feeding level and nutritional balance were the key, Dave's father Phill started using maize silage to complement pasture in the early 1990's. Today, the farm feeds around 1 tDM maize silage per cow.

Each year, Dave purchases 12 ha of maize silage from local contractor John Austin Ltd's lease block across the road. After maize silage harvest, the entire block (28 ha) is planted into annual ryegrass and Dave subleases it to winter the cows and to make grass silage.

A further 10 ha of maize is grown on farm under a strip-till system. This season half of the home crop has been grown on the effluent area with very low fertiliser inputs.



THIS PAGE LEFT TO RIGHT Phill Swney as he appeared in the June 1994 Dairy Exporter; Phill at the stack, 31 years later.

"Using a combination of growing maize silage on farm and buying it in gives us all the feed we need while diluting the average cost".

The Swneys have grown Pioneer® brand maize hybrids for more than 30 years and appreciate the longterm relationships they have with their local Pioneer representatives as well as John Austin over that time.

When it comes to maize silage, they are looking for silage quality as well as a reliable yield.

"We take a sample out of every stack and carefully analyse the results" says Dave. "A key goal is to maximise the starch content of our maize silage and there is a significant difference between hybrids. We like the quality of the maize silage we get from P0900".

For supplementary feeding, the farm relies solely on a 220-cow feedpad and a recently purchased mixer wagon.

"The feedpad is an essential part of our farm system and the mixer wagon has been a great addition because it has allowed us to get a more even distribution of minerals" says Dave. "You have to invest in the right tools if you want to push production and profit".

Calves remain on the farm until the 1st of May and are grazed off for 12 months, returning home before their first calving at the start of July.

The main herd starts calving on the 10th of July. To minimise the risk of metabolic disease, springers are fed a maizesilage-based diet that includes palm kernel, molasses, straw, anionic minerals, and a "sniff" of pasture.





GG It's a great base feed that works extremely well with pasture and helps us maximise the amount of homegrown feed we harvest gg

"Because we are calving so early, we feed high amounts of supplementary feed to start with and slowly ease off once the grass starts growing" says Dave. "We typically don't make any pasture silage on farm; we are stocked high enough that we eat all the grass we can grow".

Cows normally peak at around 2.3 kgMS/cow on a diet that comprises maize silage, palm kernel and pasture.

The cows are run in two herds. In past seasons, the young herd has been once-a-day all season. This has helped overcome historic lameness issues due in part to the large walking distance on a long, skinny farm.

"The furthest paddock is 2.2 km from the cow shed" says Dave. "It burns up a lot of energy walking that far twice daily".

This season, both herds started on twice-a-day milking, with the young herd moving to once-a-day after Christmas. The farm's summer milking schedule is unique. Cups go on at 5:30 am in the mornings, and the older herd is grazed in nearby paddocks and milked again at 1 pm. "Research has shown you don't compromise production as long as there is no more than 18 hours between milkings" says Dave. "Lunchtime milking seems to suit both us and the cows well".

In the autumn, maize silage is coupled with palm kernel and grass silage to extend the lactation and meet cow condition and pasture cover targets by drying off.

Four years ago Alice, who was formerly a Senior Catchment Officer with the Waikato Regional Council, turned the tennis court into a successful



native plant nursery trading as The Native Dairy Farmer. As well as creating non-farming income, the nursery allows her to spend more time with their sons Walter (2) and Rex (1).

Dave, who is on the Smaller Milk and Supply Herds (SMASH) Trust and National Committee, enjoys the handson aspect of a family farm and has no real desire to expand the operation. Instead, the couple plan to consolidate



their current system by focusing on efficiently increasing pasture and crop harvested and per-cow production. Adding value to the farm by further native plantings, including shade trees, is also a priority.

While his top focus is growing and harvesting as much high-quality pasture as possible, Dave believes maize silage will continue to play an important role in his future farm system, especially as the climate becomes more challenging.

"It's a great base feed that works extremely well with pasture and helps us maximise the amount of homegrown feed we harvest" says Dave. "I'd like to feed a bit more and rely less on imported feed".

Listen to an extension of this story on the Feed for Thought podcast by scanning the QR code:





Farm

- Milk 410 cow Friesian x Jersey (F8J8) herd on 123 ha (effective)
- In 23/24 season, produced 189,500 kgMS or 462 kgMS/cow
- 10 ha maize grown (P0900) on farm, 12 ha bought in
- Feed 1 tDM maize silage per cow

ABOVE LEFT Walter, Dave, Alice and Rex. ABOVE RIGHT Pioneer Area Manager Matt Towers with Phill and Dave Swney.

PIONEER® BRAND PRODUCTS


Majze

e are pleased to offer growers a powerful portfolio of high yielding, top quality silage hybrids carefully selected to meet the needs of local livestock farmers.

Our industry-leading hybrids are the result of Pioneer's world-class germplasm library, advanced conventional breeding technologies and extensive global product testing and advancement programme.

Each year Pioneer's elite genetics are tested in around 160 maize silage trials across New Zealand. Our extensive trialling programme allows us to identify superior hybrids and to position them in the growing environments where they will perform the best.

Our field team is also based in the main maize-growing regions. Give one of them, your local merchant or contractor a call for advice on choosing the right product for your paddock.





Pioneer's Optimum[®] AQUAmax[®] hybrids are highly resilient in challenging conditions and responsive to favourable ones. These hybrids are bred to include

key traits that improve your crops root system and silk emergence, among other agronomic characteristics,

AQUAmax[°]

to manage drought stress. And this season we are proud to have P0710 as the newest recruit in our lineup of high-performing Optimum[®] AQUAmax[®] hybrids. See

> page 20 for more information on this exciting new hybrid which joins P0900 and P9911, our other high-performing Optimum[®] AQUAmax[®] offerings.

Further reading on Pioneer hybrid performance

The 2025 edition of our maize silage research publication provides comprehensive data to help growers make informed decisions on which Pioneer hybrid to plant. It's packed with hybrid performance comparisons plus the results of recent agronomic and farm systems projects. To download a copy visit pioneer.co.nz/maize-silage or scan the QR code below.



SCAN THIS QR CODE TO DOWNLOAD YOUR COPY





NEW Dioneer P7179	
Silage yield	9
Grain yield	9
Staygreen 7	
Drought tolerance 6	
Early growth 7	
Stalk strength 7	
Root strength 6	
Starch and sugar	9
Whole plant digestibility 8	

Quickest option for the coolest growing regions.

Max 9

Similar in type but very much quicker to harvest than **P7364**.

- Combines superior early growth, drought tolerance and staygreen.
- Tall with low ear placement and strong standability.
- Delivers high yields of quality silage in the coolest maizegrowing regions.
- Plant at similar plant populations used for **P7364**.

An important earlier companion hybrid to **P7364** for Central Plateau, high-altitude Taranaki and high-altitude and latitude South Island growing regions.



NEW
P7364

CRM / 3			
Silage yield			9
Grain yield			9
Staygreen		8	
Drought tolerance		8	
Early growth	7		
Stalk strength	7		
Root strength	7		
Starch and sugar			9
Whole plant digestibility			9

Max 9 The new standard for yield and earliness.

Better husk cover and quicker to harvest than **P7524**.

- Dependable agronomic package with high ratings for early growth, drought tolerance and staygreen.
- Moderate in plant height with low ear placement and superior standability.
- Delivers high silage yields for maturity with superb energy and digestibility ratings.
- In medium to high-potential situations, plant 5,000 plants/ hectare more than was applied in the past for **P7524**.

Replaces **P7524** for growers requiring a hybrid earlier than **P7647** or **P8086**.



Max 9

Delivers superior yields of top-quality silage.

Expands the options for cooler maize-growing regions.

- Combines strong early growth and staygreen to deliver silage with excellent energy and digestibility ratings.
- Trials show **P7647** is slightly quicker to reach harvest drymatter and is higher yielding than **P7524** which it replaces.
- Tall for maturity, should be planted to achieve plant populations between 105,000 and 115,000 plants per hectare depending on paddock yield expectation.

An important option for South Island growers looking for a hybrid maturity between **P7364** and **P8086**.

NEW Dioneer P8086
CRM 80
Silage yield
Grain yield
Stavareen 7
Staygreen 7
Drought tolerance 7

9

Early growth Stalk strength Root strength

Starch and sugar

Whole plant digestibility	7	
Reliable early with excellent		I
valuo		

P8086 is similar in type, maturity, and management requirements to **P8000** which it replaces.

- A long cob packed with deep dent grain to produce silage with high digestibility and energy.
- Moderate in height with low ear placement, strong standability, drought tolerance, staygreen and Northern Leaf Blight ratings.
- Trials show **P8086** delivers higher silage yields than P8000.

Grow alongside P7647 or P8240 depending on maturity and disease resistance requirements.

P	
	P8240
	CRM 82
Silage yield	

0

Grain yield				9
Staygreen			8	
Drought tolerance		7		
Early growth	6			
Stalk strength	6			
Root strength		7		
Starch and sugar		7		
Whole plant digestibility		7		
			lax 9	

Bulk and energy to fill the vat.

P8240 is a very tall, high-yielding silage and grain hybrid backed by strong drought tolerance, staygreen and standability.

- · Delivers top silage yields, with superior feed quality for optimal milk production.
- A balanced agronomic package including superior roots which are a real asset in this maturity.
- Established plant populations should be matched to assessed paddock yield potential.
- Where Northern Leaf Blight is a seasonal concern, consider planting **P8086**.

Well adapted to Central Plateau, Taranaki, Lower North Island and South Island growing regions.

PIONEER. P8333	
CRM 83	
Silage yield	9
Grain yield	9
Staygreen 7	
Drought tolerance 7	
Early growth 8	
Stalk strength 6	
Root strength 5	
Starch and sugar 7	
Whole plant digestibility 7	

Max 9

Highly productive mid-maturity option.

P8333 is a tall bulky plant with a long grain-filled ear, supported by strong all-around agronomics, superior drought tolerance and staygreen.

- Delivers top silage yields for maturity, with impressive energy and digestibility.
- An important mid-maturity option between P8086 and P8666.
- · Optimum established populations are approximately 5,000 plants per hectare less than applied for **P8086**.

Yields particularly well in the South Island and in the cooler regions of the Lower North Island where this maturity is required.



PIONE		2
P866 CRM 80 Silage yield	_	8
Grain yield Staygreen	6	8
Drought tolerance		8
Early growth Stalk strength	6	8
Root strength 5 Starch and sugar	6	
Whole plant digestibility	6	

Grows well, yields very well and feeds even better.

Max 9

A bulky plant with superior early growth, drought tolerance and strong all-round agronomics.

- Produces silage with high grain content that cows will thrive on.
- Plant to establish 100,000 to 115,000 plants per hectare.

Widely adapted from Ashburton to Dargaville.

PIONEER	,
P8711	/
CRM 87	

Silage yield		8	
Grain yield			9
Staygreen		8	
Drought tolerance		8	
Early growth	7		
Stalk strength	7		
Root strength	7		
Starch and sugar			9
Whole plant digestibility			9

Yield with superb quality for Northern Regions.

Max 9

A tall hybrid with superior roots and stalks.

- High ratings for drought, Northern Leaf Blight, Rust, and staygreen deliver seasonlong silage appeal and yield stability. These all combine to support a wide harvest window.
- Produces silage with excellent energy and digestibility desired by high-productivity herds.
- Research results show **P8711** is most productive in northern regions from Northland to Hawke's Bay particularly where standability and Northern Leaf Blight are significant concerns.



PIONEER.	
P92575 CRM 92	
age yield	
ain yield	
iygreen	
ought tolerance	8
ly growth 7	

Silc

NEW

Grain yield			9
Staygreen			9
Drought tolerance		8	
Early growth	7		
Stalk strength	7		
Root strength	7		
Starch and sugar			9
Whole plant digestibility			9

Max 9

Solid, balanced hybrid with top-ofthe-line foliar health.

Plant where Northern Leaf Blight, Rust, standability and drought tolerance are seasonal concerns.

- Competitive silage yields compared to **P9400**.
- Moderately tall with strong agronomics, superior roots, and stalks.
- Delivers high and stable silage yields.
- Late season staygreen and plant health delivers a wide harvest window and silage with exceptional digestibility and energy.

Adapted to all North Island growing regions where this maturity meets growers' objectives.

S Pioneer hybrid trait comparison ratings are only valid within a range of + or - 4 CRM. These descriptions mainly feature product strengths. When choosing bubrids also rating areas for the table on page 62 Contact, your local Pioneer Area Manager or march at feature for the stable on page 62 Contact, your local Pioneer Area Manager or march at feature for the stable on page 62 Contact, your local Pioneer Area Manager or march at feature for the stable on page 62 Contact, your local Pioneer Area Manager or march at feature for the stable on page 62 Contact, your local Pioneer Area Manager or march at feature for the stable on page 62 Contact.

PIONEER P9400 CRM 94	•
Silage yield	7
Grain yield	7
Staygreen 6	
Drought tolerance	7
Early growth	7
Stalk strength	7
Root strength	7
Starch and sugar	8
Whole plant digestibility	8

Max 9

Stands tall – delivers big time.

A tall, dense plant producing high grain content silage with superior digestibility.

- Strong agronomically with a sound all-round disease resistance offering.
- When planting in early spring into cold wet soils consider planting **P92575** or **P9650**.

A popular high-yielding early choice in Northland and Waikato while giving stable yields in Taranaki and Lower North Island as a mid to full-season hybrid.



NEW DIONEER P9650

CRM 90	5		
Silage yield			9
Grain yield			9
Staygreen		7	
Drought tolerance		7	
Early growth	6		
Stalk strength		7	
Root strength	6		
Starch and sugar			9
Whole plant digestibility			9
		N	/ax 9

Security with performance.

Offers yield stability for silage and grain.

- Moderate in plant height with an erect leaf habit, strong standability and drought tolerance.
- Agronomically balanced with a strong Northern Leaf Blight resistance while producing excellent silage quality.
- Waikato research trials show P9650 was 1% wetter at harvest than P9400 but delivered 800 kgDM/ha more yield.

A useful mid-maturity option between **P92575** and **P9978** which is widely adapted to North Island growing regions.



Max 9 |

Top yielding, drought buster.

A key maturity option in the Optimum® AQUAmax® range provides growers more yield per drop – rain or shine!

- Tall, showy hybrid delivering yield stability in this maturity.
- A widely grown, imposing all-round hybrid.
- Top agronomics for reliable yields.

Where Northern Leaf Blight is a concern, consider planting **P9978** or **P0362**.



PIONE P997			
CRM 9			
Silage yield			9
Grain yield			9
Staygreen		7	
Drought tolerance		7	
Early growth	6		
Stalk strength	6		
Root strength	6		
Starch and sugar			9
Whole plant digestibility			9

Defensive. Stable. Productive.

P9978 delivers a great all-round package with superior foliar health and silage eye appeal.

Max 9

- A moderately tall plant with low ear placement, strong roots and stalks, superior drought tolerance, staygreen and Northern Leaf Blight resistance.
- Delivers top silage yields in this maturity with excellent feed quality.
- When planting early or into cold, wet soils, switch to P9650 or P9911.

Now widely grown in all North Island growing regions where this maturity is required.

		×	رو	8	
Ρ	10				l
	PO	6	4		

CRM 106			
Silage yield			
Grain yield			
Staygreen	7		
Drought tolerance	7		
Early growth		8	
Stalk strength 6			
Root strength 6			
Starch and sugar	7		
Whole plant digestibility	7		

Leaf disease champion delivering silage yield stability.

Max 9

A balanced all-round hybrid with desirable leaf disease resistances.

- Tall plant with sound standability, staygreen and drought tolerance.
- Superior resistance to Northern Leaf Blight and Rust for notable mid-to-late-season plant appeal.
- Produces silage with superior digestibility and energy content.
- Supplies yield stability in moderate to high-yield environments.

Plant with **P0362**, **P0710**, **P0900** or **P0937** depending on maturity requirements.



NEW

Max 9

Exceptional foliar health and yield stability – wet or dry!

Optimum[®] AQUAmax[®] drought qualities provide tolerance to water and heat stress during flowering and grain fill stages.

- Starts well with strong stress emergence and early growth.
- Superior staygreen and Northern Leaf Blight resistance delivering season long plant health.
- Relatively short plant with low ear placement, strong roots and stalks.
- Similar yield profile to P0900 and P0937 but higher energy and digestibility ratings.

An excellent new option to plant alongside **P0640**, **P0900** and **P0937**.

AQUAmax[•]

optimum

A YATES FAMILY BUSINESS

Р	IONEER.
	P0900
	CRM 109
Silage yield	
Grain yield	

9

Max 9

Grain yield				9
Staygreen			8	
Drought tolerance				9
Early growth		7		
Stalk strength		7		
Root strength		7		
Starch and sugar	6			
Whole plant digestibility		7		

A proven, stable, all-round hybrid.

P0900 is an exceptionally balanced hybrid that delivers yield stability and a wide harvest window.

- Dependable standability, low ear placement, Optimum[®] AQUAmax[®] drought tolerance, great foliar health and staygreen.
- Strong resistance to Northern Leaf Blight and Rust.
- East Coast growers will value superior Head Smut resistance.
- A management-responsive hybrid that will benefit from adjusting the established plant population to match yield expectations.

Extensively planted between Kaitaia and Napier alongside **P0937**.

AQUAmax

ptimum



Silage yield			9
Grain yield			9
Staygreen	7		
Drought tolerance		8	
Early growth		8	
Stalk strength 6			
Root strength	7		
Starch and sugar	7		
Whole plant digestibility		8	

Solid hybrid with great standability and foliar health.

A modern plant type with erect leaves, notable foliar health, standability and exceptional staygreen.

- Widely adapted, stable, yet high-yielding hybrid.
- Emerges strongly when planted early into cold wet soils.
- Sound Northern Leaf Blight and Rust resistances will be attractive to growers in highrisk situations.

P0937 performs best in moderate to high-yielding situations in warmer northern growing regions. Companion with **P0900** and **P1185**.

PIO		R.)	
P 1	315		
CRI	м 110		
Silage yield			
Grain yield		7	
Staygreen			8
Drought tolerance		7	
Early growth		7	
Stalk strength		7	
Root strength	6		
Starch and sugar	5		
Whole plant digesti	bility		

6

Max 9

0

Defensive from Northland to Hawke's Bay.

An imposing hybrid like **P1636** while being quicker to harvest and with much better foliar health.

- Season-long appeal is delivered by notable drought tolerance, staygreen and superior Northern Leaf Blight and Rust resistances.
- Tall with superior root and stalk strength.
- Trials show **P1315** delivers higher silage yields than **P0900** and **P0937**.
- Where Head Smut is a concern, plant **P0900**.

Companion with **P0900, P0937** or **P1636**.

NEW
PIONEER.
P1185
CRM 111

Slidge yield		
Grain yield		9
Staygreen		9
Drought tolerance	7	
Early growth	7	
Stalk strength		8
Root strength	7	
Starch and sugar	7	
Whole plant digestibility		8

Max 9

Foliar health champion delivering top quality silage.

Similar in type and management requirements to **P0900** and **P0937**.

- The standout performer in trials for Northern Leaf Blight and Rust resistances, standability, staygreen and season-long plant health.
- Delivers higher silage yields and a wider harvest window than P0937.

Plant from Kaitaia to Napier as a companion to **P0900** and **P0937**, particularly where there's been significant Northern Leaf Blight pressure in recent seasons.



PIONEER P1636

CRM 112

Silage yield				
Grain yield		7		
Staygreen		7		
Drought tolerance		7		
Early growth		7		
Stalk strength	6			
Root strength			8	
Starch and sugar	6			
Whole plant digestibility	6			

Max 9

Enjoy the agronomics of this top-yielding hybrid.

P1636 is a tall full-maturity hybrid that consistently delivers top-end yields in this key late-maturity segment.

- Long cob to produce high grain content silage.
- Combines impressive
 agronomics, drought tolerance
 and staygreen that together
 provide a wide harvest
 window.
- Plant early to maximise yields.
- In high-risk Northern Leaf Blight situations, consider P1185, P1315 or P1837.

P1636 is well adapted to all warmer northern growing regions.

PIONEER PIB337 CRM 115 Silage yield 9 Grain yield 7 Staygreen 9 Drought tolerance 8 Early growth 7 Stalk strength 6 Root strength 6 Starch and sugar 5 Whole plant digestibility

Max 9

Defensive full-season giant.

Very tall, with superior standability and resistance to Northern Leaf Blight and Rust.

- Excellent drought tolerance and staygreen which support season-long "silage eye appeal".
- Better foliar health than P1636 while being slightly later in maturity.
- This huge plant only requires established plant populations of 70,000 to 90,000 plants per hectare to optimise yield performance.

A widely adapted, stable, fullseason hybrid. Plant early in high-potential paddocks in warm northern production areas.

Companion with **P1315** or **P1636**.





CRM 103





CRM 114



Seed beatments





Selected for our genetics

Protection today, performance tomorrow

For Pioneer offers industry-leading protection for your Pioneer® brand seed.

LumiGEN[®] seed treatments protect your investment in Pioneer seed from earlyseason disease, insects, and bird damage, helping to maximise yield potential.


Proven in the field with our genetics

Verified on

our genetics

MAIZE FOR SILAGE 2025/2026 25


INSIDE PIONEER



Understanding seed challenges and how to protect your crop

Last season, some growers faced higher-than-normal challenges with pests such as greasy cutworm, Argentine stem weevil (ASW), birds and rats. While LumiGEN® seed treatments provide industry-leading protection against most early-season threats, they are not a complete solution on their own. Successful pest management requires a combination of the right seed treatment and other proactive management strategies such as fallow periods, monitoring and baiting. Taking these extra steps can make all the difference in protecting your crop's potential.

Talk to your local Pioneer representative, merchant or contractor to determine the best LumiGEN[®] seed treatment option for your growing environment.



* Research data shows the insecticide in L-400 and L-400+ is more effective at reducing plant loss due to Argentine stem weevil damage than the insecticide in L-300 and L-300+

¹ L-400 and L-400+ contains Bacillus spp which suppresses nematode damage in maize



Greasy cutworm

Greasy cutworm damages maize by cutting seedlings at ground level with most losses occurring during crop emergence. The presence of weeds at planting is a major contributor to cutworm infestations because weedy patches allow caterpillars to survive and carry over until maize plants emerge.

To control greasy cutworm, a 5-6 week weed-free fallow period before planting is ideal, though not always practical.



LumiGEN® L400 seed treatment offers short-term control under low to moderate cutworm pressure. However, if infestation levels are high, a spray application of insecticide may be necessary to prevent further damage. These insecticides are cost-effective and can significantly reduce the impact of cutworm.

Argentine stem weevil (ASW)

Maize can be at risk of damage by ASW larvae when planted less than four weeks after cultivation from pasture, annual ryegrass, or cereal crops. Partly developed larvae transfer from decaying grass tillers and tunnel into the young maize plants, which can result in:

- **Pre-emergence damage:** seedlings killed before emergence.
- Post-emergence stress: affected plants display blue-grey discolouration, and their centre leaves pull out easily. A small, round hole at the base of the plant indicates ASW entry or exit.



ASW damage typically occurs within the first four weeks of maize growth. The best way to reduce the risk is by planting maize seed treated with **LumiGEN® L400 seed treatment**. In high-pressure situations, a 4-6-week fallow period before planting can help reduce infestation levels.

Even if you have used insecticide treated seed, you should monitor your crop daily. Currently, no insecticides are registered in New Zealand to control Argentine stem weevil in a germinated maize crop. Call your local Pioneer Area Manager or merchant representative if you notice ASW damage in your maize crop.

Birds

For a reliable, safe and non-lethal way to protect maize seed from birds, consider using **LumiGEN® L-200+**, **L-300+ or L-400+ seed treatments**, which contain a scientifically formulated bird repellent. This unique formulation is based on an organic chemical naturally found in several plants, including aloe vera and rhubarb, and has been proven to be highly effective in deterring birds from eating maize seed.



Rats

Rat damage occurs typically near the paddock's edge, and there is a characteristic "hoof" shaped hole where the rat dug for the seed. After consuming a kernel, rats often continue along the row, causing significant damage. Unfortunately, seed treatments do not deter rats.

To minimise rat damage, control measures should begin in autumn or winter:

- **Baiting:** place bait stations around the field perimeter and replenish bait regularly to maintain effectiveness.
- Habitat management: reduce cover for rats by clearing overgrown weedy areas and managing waterways along the paddock edges.

By taking these steps, you can significantly reduce the risk of rat damage and protect your maize crop.



LumiGEN[®] seed treatments play a critical role in protecting your maize, but they should be part of a broader pest management strategy. Combining seed treatment with proactive control methods will give your crop the best possible start.



Looking after/ your marge seed

he highest-yielding maize crops are harvested from proven maize hybrids planted in even plant stands that have uniform emergence. Seed quality is paramount to your maize crop's success. In this article, we outline the factors that impact

maize seed quality and detail some of the steps we take to help ensure the Pioneer[®] brand maize seed you plant is of the highest quality. We also discuss what you can do to ensure seed quality is maximised right through to when you plant it in the ground.



Maize seed – a living organism

A seed is a living organism that contains the genetic material, nutrients, and energy needed to grow a new plant. A maize seed consists of three main parts: the germ, the endosperm, and the pericarp.

pericarp

endosperm

germ

The **germ** contains a living embryo which grows into a maize plant.

The **endosperm** is the starchy tissue surrounding the germ, providing energy and nutrients for growth.

The **pericarp** or seed coat protects the seed until conditions are favourable for germination.

Maize seed germination and growth can be impacted by both physical and physiological factors:

Physical damage (e.g. cracks or broken kernels) can directly damage the germ or disrupt the protective pericarp allowing moisture or pathogens into the seed.

Physiological damage, which impacts the dormant embryo, can occur when maize seeds are exposed to undesirable temperatures, humidity or too much light.

Determining and defining seed quality

Samples of all lots of Pioneer maize seed undergo a range of tests and checks to ensure they meet Pioneer's stringent quality standards.





 Uniformity of seed size and shape, and seed treatment integrity. Freedom from

contaminants such as weed seeds and inert material.



Freedom from fungal diseases.
Absence of insect pests.





Handle the bags gently to avoid cracking seeds or rubbing seed treatment off them.



Tips for looking after your maize seed

We asked Seed Quality Manager Mark Robertson how growers should look after their maize seed on farm.

Handle your seed carefully and as little as possible. If your seed arrives on a pallet, try to keep it there until planting. If you need to move the seed, handle the bags gently to avoid cracking seeds or rubbing seed treatment off them. Never throw seed bags.



Keep the seed in a cool, dry place away from direct sunlight. This will help maintain the life of your seed.



Don't store maize bags alongside chemicals which may release fumes that could impact the seed.



Control rodents and birds, which could make holes in the bags and damage the seed.

Producing high-quality maize seed

Our focus on seed quality starts in the field, continues right through the seed production process and does not end until every bag of Pioneer maize seed is planted in the ground. We asked Seed Production Manager Andrew Powell, to outline some of the things the Production Team do to ensure the Pioneer[®] brand maize seed you receive is of the highest quality.





CROP MANAGEMENT

A healthy seed crop is the starting point for high-quality seed production. Careful attention is paid to crop nutrition and the control of weeds, insect pests and fungal diseases.

DRYING

It is important that cobs are dried as soon after harvest as possible to avoid the risk of sweating. After husking the cobs are carefully lowered into the driers. Air temperature is carefully monitored and altered throughout the drying process to ensure cobs are dried at the correct rate. The final moisture content is also important because if the seed is too damp, it will deteriorate at a faster rate. Once dried to the correct final moisture, the seed quality is stabilised and the remaining processes are less time-critical.

HARVESTING

It is important to harvest at the right time. If the seed is too wet, a lot of moisture has to be removed, and this can impact its quality. Conversely, if the seed is too dry, there is a higher risk of physical damage during harvesting and processing. Seed maize is harvested on the cob using specialised equipment. The husk is left on the cob to help protect the seed throughout the harvesting process.



PROCESSING

The seed must be shelled (taken off the cobs), cleaned, sized, treated and bagged carefully to avoid any physical damage. A combination of easy let-down ladders and specially designed conveyors ensures the seed is never dropped onto hard surfaces.

TESTING

The seed testing regime is an important part of ensuring seed meets Pioneer's rigorous quality standards. Samples are routinely collected throughout the seed production process and shipped to an International Seed Testing Authority (ISTA) accredited laboratory for testing. Every batch of maize seed that leaves our Gisborne production plant has been tested for genetic purity, germination and vigour.


PIQNEC

STORAGE

Heat and moisture are the biggest enemies of seed quality. Pioneer seed is stored under controlled temperature (10°C) and humidity (50%) conditions which slow down seed metabolic processes helping to maintain seed quality.

FARM SUCCESS STORIES

Maize helps power production



mix of foresight and innovation led the Singh family from a traditional dairy farm to a 1000-cow freestall barn system.

Lawwal Holsteins represents the vision of the late Aman Singh whose aspirations were to feed cows to their potential, increase production per cow and look after the environment, while utilising the land in a better way.

Today, the farm is managed by his three children, Arjun, Anjena, and Amreeta, and owned by the family, including their grandparents, Karamijit and Mindy, and their mother, Daljit.

In the 2023-24 season the 250 ha farm, which is located at Whitikahu near Hamilton, milked 800 cows, producing an impressive 600,000 kgMS (750 kgMS/cow or 2,400 kgMS/ha).

LEFT Arjun Singh chats with Pioneer Regional Manager Brent Bishop.

FARM SUCCESS STORIES

The current system of farming 650 kg Holstein cows indoors year-round represents an evolution of cows as well as farm system. The original Law Road property of 116 hectares was purchased in 1989 and the family ran 270 cows in two herds (one Jersey, the other Holstein Friesian) through a 22bale rotary shed.

The next two decades saw the dispersal of the Jersey herd, an increase in cow numbers and the installation of a feedpad and a 50-bale rotary. The purchase of an adjoining property in 2013 brought the farm to its current size.

The next year the family debated their next move, tossing up the alternatives of buying more land or moving to a different production system. Aman travelled to America, Europe and the South Island to view systems and capture the best ideas.

The culmination of his trip was the family's decision to build a 1000-cow freestall barn adjacent to the existing cow shed. A number of different designs were studied, and a final design that suited the climate and maximised cow comfort was chosen. The 1000-cow freestall barn was commissioned in April 2018.

The 245 m long barn is split into quarters and the farm runs four milking herds. Cows are separated based on their stage of lactation and production level and each herd gets a different total mixed ration (TMR) suited to their unique nutritional needs. Maize and grass silage provide the base for the ration which is then balanced with bought in protein, starch and energy sources as well as minerals.

Seventy-five percent of the TMR is homegrown. Each season, the family plants 200 ha of Pioneer® brand maize for silage. The remaining 50 hectares are used for youngstock grazing, allowing the family to operate a closed system.

In the 2024-25 growing season, they have planted a mix of Pioneer maize hybrids, including P8240, P9400, P9978 and P0937.

"We are looking for hybrids which deliver a high grain yield because we want to maximise metabolisable energy and starch intake" says Arjun. "Small variances in this can have a large impact on per cow production. Good disease resistance and drought tolerance are also important".

The last few Waikato summers have delivered several cyclones, so good standability ratings are now also important.

"We choose to plant Pioneer because the service we get from our local Regional Manager Brent Bishop is second to none" says Arjun. "The replant policy is another selling point for us; we hope we don't need to use it, but having it there gives us peace of mind".



ABOVE The Singhs' 1000-cow, 245 m long freestall barn. **RIGHT** Arjun Singh

The family are ardent supporters of the Rural Support Trust and Arjun says Pioneer's contribution to this valuable organisation "shows a lot about their character".

Maize crops are planted with starter fertiliser only and the farm relies on the use of effluent (both liquid and solids) which is injected 200-300 mm into the ground for the remainder of the crop's nutrients with the exception of sulphur. The target average maize silage yield is 25 tDM/ha, which brings the cost to 13c/kgDM in the stack.




After the maize silage is harvested, the effluent is applied via a dribble bar. Then, the land is direct drilled into annual ryegrass and harvested three times for grass silage before returning to maize.

"We get 7-8 tDM/ha from the ryegrass, giving a total annual drymatter yield of around 32 tDM/ha" says Arjun. "It's a lot more than we could grow under any other pasture or cropping system".

The farm's careful nutrient management, which includes twiceyearly soil testing, and lack of synthetic fertiliser contributed to the farm being named the Regional Supreme Winner of the Waikato Ballance Farm Environment Awards in 2023.

While the business focuses on maximising milk production, selling embryos and livestock provides a significant secondary income stream for the farm. Top cows are flushed, and embryos are transferred to recipient cows. All replacements are genomic tested at birth, and this provides ratings for a large number of traits including milk fat and protein percentage, methane production, feed conversion efficiency and net genetic merit.



Last year the family sold a half share of the highest conformation heifer in the

and have stood the test of time".

So, what does the future look like?

Our aim is to have 1,000 cows in milk year-round and to increase cow production by improving genetics and further refining our TMR system" says

GG The replant policy is another selling point for us, we hope we don't need to use it but having it there gives us peace of mind \Im

world to an American buyer. The heifer "Bellarina" remains on the farm while her embryos are being flushed and shipped back to the U.S.A.

Cows are all artificially inseminated with a high use of sexed semen. Seventy percent calve in February and the balance in September.

"About 90% of the herd carry North American genetics, and we utilise Semex and World Wide Sires bulls, says Arjun. "These cows are best suited to our production system; they efficiently produce high milk volumes Arjun. We have recently started a new partnership to bring Showbox Sires to New Zealand from North America.

"The weather extremes we are currently experiencing will continue to make it even more attractive to grow maize, and we would like to have more maize silage on hand so we can carry a buffer from season to season".





Farm

- Milk 800 Holstein cows indoors year-round on 250 ha
- In 23-24 season produced 600,000 kgMS or 750 kgMS/cow
- 200 ha maize grown averaging 25 tDM/ha
- Grows P8240, P9400, P9978 and P0937.






INSIDE PIONEER




Pioneer





in NZ











A YATES FAMILY BUSINESS 38

÷





























INSIDE PIONEER

his season we celebrate 50 years since the signing of the seed production and distribution agreement between Pioneer Hi-Bred International Incorporated and Philip Yates, who at the time was the Managing Director of Arthur Yates & Co. Ltd and subsequently the Founder of Genetic Technologies Ltd, another Yates family business.

The signing marked the first step in the launch of Pioneer® brand maize and other products in New Zealand and the start of an enduring relationship between Pioneer and the Yates family, both of whom have had a long history in the global seed business. Pioneer Hi-Bred International was founded by Henry A. Wallace who later in life was elected Vice President of the USA under President Franklin Delano Roosevelt. Henry was one of a handful of people in the world who initially recognised the immense opportunities that could be gained by growing hybrid maize. He began experimenting with the crop in high school and while at Iowa State College, became fascinated with the relatively new science of hybrid genetics. After graduating in

On Tuesday 18 December 1975 in Des Moines, Iowa, Philip Yates (second from left) and the Pioneer executives sign an agreement for Arthur Yates and Co. Ltd to be the New Zealand distributor and producer of Pioneer* brand seed.



INSIDE PIONEER

1910, Henry started breeding hybrid maize and by 1923 had produced a high-yielding hybrid he called Copper Cross. In 1924, it became the first hybrid to win the gold medal in the lowa State Corn Yield Contest. In 1926 Wallace incorporated the Hi-Bred Corn Company, which later became Pioneer Hi-Bred International.

A name synonymous with seeds

The Yates family's involvement in the seed industry started in England in the late 1700's. James Yates, a cotton importer purchased a small quantity of Egyptian cotton seed and sold it to the colonist farmers in the southern part of the present-day U.S.A. This proved an inspired move as the demand for cotton seed grew and very soon James relinquished his cotton importing business and became a seed merchant.

Almost a century later, James' great grandson Arthur who was

an asthmatic, took a sea voyage away from the smoky cold air of the industrialising British Midlands to New Zealand. Arthur worked as a shepherd and scythed grass seed from roadsides to supplement his income. The seed revenue quickly outstripped his shepherd's wages, and in 1882 Arthur moved to Auckland, rented a rickety wooden shop in Victoria St West and opened a specialised seed business.

Philip Yates takes the reins


Arthur's younger brother Ernest Yates travelled from England to join him in business in 1886. Ernest was followed in the trade by his son Norman and in 1973 grandson Philip became Chief Executive and Managing Director.


Arthur Yates and Co. Ltd prospered and Yates' Reliable Seeds became a household name, wholesaling seed of every kind from flowers to vegetables, broad acre agricultural seeds to bird seed. While their product range was large, Philip was always looking for superior seed products. A magazine article about hybrid wheat caught his interest and ultimately led to a phone call to Pioneer Hi-Bred's Overseas Division based in Des Moines, Iowa.

Signing with Pioneer

After a whirlwind tour of Pioneer's U.S. operations and breeding programmes, Philip came to realise there was more near-term potential for hybrid maize than hybrid wheat in New Zealand.

Philip was impressed by Pioneer's significant library of elite maize germplasm, their commitment and innovation in plant breeding, and above all else, their conservative family values and customer-focused "Long Look" philosophies. On Tuesday 18 December 1975, an agreement was signed in Des Moines, Iowa and the Yates Company became the New Zealand distributor and producer of Pioneer[®] brand seed.





The timing of the relationship between the two companies could not have been better. In the 1970's Pioneer invested significantly in developing hybrids with ever-higher yields. Their new maize hybrids were breaking U.S.A. yield records and sales were on a rapid upward trajectory increasing five-fold from 1972 to 1980.

A slow process

While Pioneer had a range of exciting new products, New Zealand's tight biosecurity requirements meant that only 48 individual maize seeds could be imported at a time. On arrival, seed had to be treated with a toxic combination of fungicides which meant only half would remain viable. Getting to the point of commercial release of a Pioneer hybrid was therefore a slow and labourintensive process. "We started with a handful of Pioneer parent seed which we had to multiply before we could get enough to produce even trial quantities of seed" says Philip. "It took a very long time to multiply the quantity of inbred seed to produce commercial volumes".

The first Pioneer maize hybrid to be commercialised in New Zealand was 3709. The next hybrid was 3591 and a couple of seasons later 3901 was released as a shorter maturity option for lower North Island growers.

A winning hybrid takes off

Winning an independent maize grain trial at Kaipaki in the western Waikato was a turning point for sales with the two Pioneer hybrid entries 3709 and an experimental outstripping all the commercially available competitor hybrids. "Pioneer maize seed wasn't hard to sell" according to Richard Jellie who was a Waikato-based Sales Manager for Yates in the 1970's. "3709 was a reliable hybrid which produced higher yields than longer maturity competitor hybrids and it didn't fall over. Local growers had heard of the brand's success in the U.S. market and couldn't wait to get their hands on it".

The improved standability was a real bonus because prior to 3709's introduction it was not uncommon for people to walk alongside the combine standing up plants so the cobs could be harvested.

Early days of seed processing

The initial Pioneer seed crops were hand-picked and sun dried, but as sales grew Arthur Yates and Co. Ltd built a seed production plant at Waharoa in the eastern Waikato. Cobs were sorted and dried and the seed was then sent to the Manawatu for conditioning.

Neville Westbury, who started with Yates in 1978, can remember the challenges of drying seed maize in the damp Waikato climate.

"Maize was planted a lot later and the seed harvest didn't take place until June or July" says Neville. "When it was foggy you could stoke the furnace all night and you were lucky if the seed hadn't gained moisture by the morning".

The seed drier was fired by burning the cob cores as well as coal. This saved fuel costs but ultimately proved to be a costly decision.

"The level of technology was low at the time and the smoke and gases discharged from incomplete cob combustion was so acidic that after a short few years the new seed production plant was almost completely rusted away" says Philip.

An end and a beginning

As the maize seed production plant was coming to an untimely end, so too was the Yates family's involvement in the Yates company. In 1985, Equiticorp, a recently established investment bank, took control of the company via a share market raid and at the age of 53, Philip was dismissed and given two days to vacate his office.

LEFT Philip Yates as he appeared on the front cover of The New Zealand Financial Times (July 1979).

ABOVE An advertisement for P3709, the first Pioneer hybrid commercialised in New Zealand.



Pioneer executives in the U.S. were unimpressed by the turn of events which did not sit well with their business ethics. They cancelled their distribution agreement with the now Equiticorp-controlled Yates company and offered Philip the Pioneer representation in New Zealand. Genetic Technologies Ltd was formed.

The new company had humble beginnings with the original two staff members, Operations Manager David Lobb and Administration Assistant Patricia Kessler, sharing an office with Philip in the downstairs of his Auckland home.

GG Our company vision is to enhance the future of New Zealand agriculture gg

ABOVE Will and Philip Yates.

Ideal conditions for growth

For the first three seasons, Genetic Technologies imported Pioneer® brand maize seed or had it produced locally under contract. In 1989 Philip employed Phil Evans as Production Manager and in 1990 a new seed production plant was commissioned in Gisborne.

"The growing conditions in the Gisborne region were ideal for maize allowing us to produce some of the world's finest quality seed" says Philip.

Over the next few years, Genetic Technologies invested significantly in hybrid and agronomic research as well as continuing to promote the adoption of maize silage into the dairy industry. The field team grew to include Area Managers located in all maize growing areas as well as Dairy Specialists and a sizeable research team.

"We quickly realised it was impossible to get high quality trial data from a handful of trials" says Philip. "Pioneer had an enormous pool of maize genetics and we needed to invest in a large-scale local hybrid evaluation programme which would allow us to accurately identify those which performed best under New Zealand's wide range of growing conditions".

Vision for the future

Philip's son Will joined the company in 1990 and became the seventh generation of the family to be involved in the seed trade. Will "did his time" working alongside the field team including in the South Island market. Today he leads the management team for Genetic Technologies Ltd in New Zealand.

"Our company vision is to enhance the future of New Zealand agriculture, and we are investing in a range of hybrid, agronomic, farm systems and environmental research which will help local growers and users of Pioneer[®] brand products build profitable and sustainable businesses" says Will.

"It is a real privilege for the Yates family to have represented Pioneer in the New Zealand market for the past 50 years and I want to personally thank all the farmers who have planted Pioneer as well as the many merchants, contractors and Pioneer team who have supported them over all those years".



<section-header><section-header>

STONEY

ab 50



SCAN HERE TO LET US KNOW YOUR FLEECE SIZE. To celebrate 50 years of Pioneer in New Zealand, we're giving away a **limited-edition Stoney Creek fleece** with every maize seed order placed before 31 July 2025.* Plus you'll get our **100% Seed Replant Benefit Cover***, meaning you'll receive replacement seed at no charge if a crop needs replanting.

Talk to your Pioneer Area Manager or local merchant | www.pioneer.co.nz/early-order



*One fleece per customer. Terms & conditions apply. See www.pioneer.co.nz/early-order

Delivering more higher-quality silage

elivering fast, efficient fermentation to protect nutrients, improve drymatter recovery and improve aerobic stability, Pioneer[®] brand inoculants maximise the quality of your silage from harvest to feeding.

Technology that exceeds just a bottle

to

ONEER

e inoculan

With a range of product options suitable to be used across a number of crops and a field team to ensure you get the best outcome, Pioneer inoculants

help deliver better silage for your farm every season. There are many reasons Pioneer inoculants maximise your silage returns:

Patented bacterial

strains. Each bottle of Pioneer® brand inoculant contains patented strains selected from Pioneer's world-leading collection of naturally occurring silage fermentation bacteria.

Crop-specific inoculants. Research has shown that some bacterial strains perform very well in one crop but not as well in another. For this reason, Pioneer[®] brand inoculants consist of individual strains and combinations selected to optimise the fermentation of specific crops.

> Rapid React[®] aerobic stability technology. Pioneer[®] brand inoculants with Rapid React[®]

technology create stable feed in just

seven days, meaning you can feed your silage one week after harvest whilst still capturing the benefits of a cool bunker face, reduced drymatter losses and extended bunk life.

Local and overseas product research¹.

Pioneer[®] brand inoculants have been tested and proven under a range of ensiling conditions.

We can provide trial data for each of our quality inoculant products.

Quality assured with

ISO 9001:2000. All Pioneer inoculants are produced and managed to the highest quality control standards. We publish guaranteed bacteria levels on the label of every bottle.

Comprehensive in-field support.

Every bottle of Pioneer[®] brand inoculant is supported by an experienced, local field team with extensive silage-making expertise.



¹Trial results available on request.

Shaun Body Pioneer® brand products Portfolio Manager

According to Shaun

Shaun Body has seen firsthand the difference that feeding quality silage can make to livestock performance. He believes local farmers increase the return they get from feeding silage by applying best management harvest, storage, and feed-out practices, including applying a proven Pioneer[®] brand inoculant.

"Often there is little difference between the cost of high-quality and poor-quality silage, but the difference in milk or meat producing potential can be huge" says Shaun. "It's important to do the job right so that you maximise the return from your silage".

"Part of 'doing the job right' is ensuring you get the best fermentation possible, and that means applying efficient fermentation bacteria rather than relying on the naturally occurring strains".




NEW 11C33 with Rapid React[®] aerobic stability technology


Pioneer[®] brand 11C33RR maize silage inoculant has been proven to reduce heating and feed-out losses under New Zealand silage-making conditions. It improves silage quality by providing a low terminal pH and a desirable volatile fatty acid profile. New Zealand trials show that Pioneer's advanced Rapid React[®] aerobic stability technology creates stable feed in as little as 7 days².

While 11C33RR is supported by overseas data, New Zealand's warm, humid growing and feed-out conditions can result in silage which is more prone to heating. For this reason, we tested four local maize silages that had been inoculated with 11C33RR. These were opened early (day 9-11) or late (day 60) and compared to an untreated control³. The results showed the 11C33RR improved aerobic stability by 43.2 hours for early opening and 97.3 hours for late opening.

Improved aerobic stability and reduced heating



Inoculating with 11C33RR resulted in 43.2 more hours before heating and 4.2% less DM loss when compared to the untreated control when the stack was opened at day 9-11.




Inoculating with 11C33RR resulted in 97.3 more hours before heating and 6.3% less DM loss when compared to the untreated control when the stack was opened at day 60.





Pioneer[®] brand 11C33RR

Maize silage-specific inoculant with next-generation L. buchneri designed to reduce heating and improve silage quality. Rapid React aerobic stability technology provides less heating and stable feed in 7 days².





Improves drymatter recovery and silage quality of all forage crops.

² Improved aerobic stability and reduced heating are relative to untreated silage. Actual results may vary. The effect of any silage inoculant is dependent upon management at harvest, storage and feed out. Factors such as moisture, maturity, chop length, and compaction will determine inoculant efficacy. ³ 2022 NZ inoculant study; using the Honig method to determine heating and DM losses.



Pasture and other crop inoculants



Pioneer[®] brand

11G22RR

A grass/lucerne/cereal silage inoculant with next-generation L. buchneri designed to reduce heating and improve silage quality.



Pioneer[®] brand

Lucerne-specific silage inoculant selected to best use plant-available sugars, maximising fermentation quality, silage digestibility and animal performance.





Produces top-quality pasture silage with enhanced fermentation for highproducing dairy cows and specialised beef production¹.





Improves drymatter recovery and silage quality of all forage crops.

¹ Trial results available on request.

Pioneer brand Summer feed

hen it comes to summer feed, forage sorghum and sudan grass hybrids are a great option if you are in a warm part of the country. When looking to plant in November or December, choosing forage sorghum and sudan gives you a high-yielding, drought-tolerant option that can be grazed or harvested and stored as silage or hay for when you need it most.

Drought tolerant

Forage sorghum and sudan grass hybrids have an extensive secondary root system and a waxy leaf surface giving them a significantly higher water use efficiency than most pasture species. This makes them the ideal crop for warm, dry weather. If you are after a cropping option that can tolerate dry conditions, look to plant some Bettagraze or SSS this season.



9
9
9
9
9

Max 9

The next generation hybrid.

Super Sweet Sudan (SSS) hybrid is quick to graze and sustains multiple grazings. Fine leaves make SSS suitable for grazing with sheep or cattle or making high quality baled silage or hay.

Key features:

East food

- Sudan x sudan grass
- Quick regrowth allows multiple cuts or grazings
- Prolific tillering habit and superfine stems
- Sweet and leafy for enhanced palatability

Bag size: 15 kg

Recommended planting rate: 15-25 kg/ha

Planting depth: 2.5-3.5 cm

	mann				
Fast feed					9
Silage making					9
Hay making				8	
Sheep grazing		2.5	7		
Beef grazing				8	
Dairy grazing					9
				Max	9

 $\langle \mathbf{a} \rangle$

PIONEER

TTAGRA7

First to plant, first to feed.

Excellent recovery from grazing or cutting, along with delayed flowering makes Bettagraze a versatile, easy to manage summer feed option. Excellent palatability and a high leaf-to-stem ratio.

Key features:

- Sorghum x sudan grass
- Fine stems and disease-free leaves
- Larger seed size for better establishment in dry conditions.
- Suitable for grazing with dairy or beef cattle.

Bag size: 25 kg

Recommended planting rate: 25-45 kg/ha

Planting depth: 3-5 cm

Scan for more information regarding Pioneer summer forage hybrids:



Key

1 = Poor, 9 = Excellent. Ratings based on Pioneer research comparisons with other Pioneer[®] brand sorghum/sudan hybrids.

A proud history, an exciting future

long family farming history has made a significant contribution to Will and Louise Bailey's dairy farming journey.

The couple, who have two children – Stella (5) and Blake (almost 2) – have been contract milking 650 Holstein Friesian cows at Swannanoa in North Canterbury for the past seven seasons. Their 225 ha property consists of a 185 ha milking platform and a 40 ha run-off used to grow maize silage as well as kale for cow wintering. The couple co-owns the farm in partnership with Louise's family and Will's grandparents.

Will and Louise's Knightlea Stud combines Knighton, the name of Louise's family's beef and cropping farm and Karalea, the name of Will's grandparents' original Holstein-Friesian stud.

"There is a lot of history and breeding in the herd" says Will. "Grandad registered his first Holstein Friesian cow in 1965. I'd like to carry that herd on for another generation".



When it comes to breeding, the focus is on creating a high-producing cow that's not too big, has good conformation, good feet and legs, excellent fertility and an all-round correct cow.

The herd is split calving with approximately 500 spring (1st of August) and 150 autumn (1st of March) cows.

"We split calve because we have a winter milk contract to fulfill" says Will, "High-producing Holstein-Friesians can be a bit trickier to get in calf, so having two chances is another bonus of the system".



FARM SUCCESS STORIES





- Milk 650 Holstein-Friesian cows on 225 ha (185 ha milking platform, 40 ha run-off)
- In 2023-24, 630 cows produced 375,000 kgMS or 595 kgMS/cow
- Planted 8 ha maize and will purchase in a further 17 ha maize silage
- Last year P7524 yielded 23 tDM/ha with paddocks regrassed before winter

The mating period is 12 weeks in the spring and six weeks in the autumn. Sexed semen is used over the topperforming cows while the rest of the herd is mated to beef genetics.

In the 2023-24 season, the herd peak milked 630 cows to produce 375,000 kgMS, which equates to 595 kgMS/ cow or 1,667 kgMS/ha.

"Production is a major focus for us, and we are targeting 600 kgMS per cow each season" says Will. "Breeding is important but to get the most from our 600 kg cows we also need to focus on feeding them well".

The couple are strong advocates for maize silage.

"I worked for a contractor harvesting maize silage about 12 years ago and saw the high yield potential of the crop" says Will. "We love maize silage and wouldn't farm without it". This season they have planted 8 ha on the milking platform, and will purchase in a further 17 ha.

"We've always planted Pioneer® brand maize hybrids, and we like the shorter maturity ones best" says Will. "They deliver excellent yields of high grain content maize silage".

Will plants the farm's maize crop each year, and he also does some contract planting in the local area.

In the 2024-25 season the couple planted P7524. Last season the same hybrid grown on the milking platform yielded 23 tDM/ha and was harvested in mid-March, which was early enough to get the paddocks regrassed prior to the winter. All of the maize silage is inoculated with Pioneer[®] brand 11C33 Rapid React which helps to reduce heating and spoilage at feed out time. Historically, all the farm's silages were fed in the paddock. Pioneer Farm System Specialist Matt Dalley and Area Manager Paige Farrell helped the couple assess the financial feasibility of building a feedpad. The numbers stacked up and the pad was commissioned in July 2024.

Already, the couple are seeing significant benefits.

"Building the feedpad has allowed us to minimise silage wastage as well as pasture damage" says Will. "We've also found we are spending a lot less time on a tractor feeding out".

Spring-calving cows are wintered on kale and grass silage on the run-off. They return to the milking platform as springers and are fed a maize silagebased transition cow diet, which helps to minimise milk fever.

GG All of the maize silage is inoculated with Pioneer[®] brand 11C33 Rapid React which helps to reduce heating and spoilage at feed out time $\Im \Im$





Once they calve, cows are fed wheat, a protein meal and minerals through the in-shed feeding system.

"We don't believe there is much value in feeding palm kernel to our highproducing cows" says Will. "We would rather spend a bit more money and purchase feeds with much higher energy or protein levels". Maize silage is fed on the feedpad at about 6 kgDM per cow per day in the spring and dialled back to 2 kgDM/ cow as the pasture grows. Typically, the only months the farm doesn't feed maize silage are November, December and January.

"We aim to carry a buffer of maize silage so that we can start feeding it again in February before the next crop is harvested". While Will can see the benefits of feeding maize silage at other times of the year, he feels he gets his biggest "bang for buck" in the autumn.

"We have a lot of sloppy wet grass and maize silage helps to balance the diet, increase energy intakes and improve cow condition".

OPPOSITE TOP LEFT Pioneer Area Manager Paige Farrell and Will Bailey. **ABOVE** Will and Louise Bailey's crop of Pioneer P7524 maize.

Maize silage



Growing and Harvest Cost Guide

The costs to grow, harvest and store maize silage are estimates only based on a sample of contractor rates, typical industry charges and product costs. All costs exclude GST and are indicative as at 31 March 2025. Given the volatility of farm input prices over the last few months, we encourage you to complete your own budget prior to the start of the growing season.

To use this cost guide, enter your own growing and harvesting costs in the "My Costs" column. For help and notes on this table refer to the assumptions on the next page.

	5-26 Average Estimated Cost	Average estimated		
exclu				
èro	wing Costs	Typical fertility	High fertility	My farm (\$/ha)
	Cost of leased land ¹	?	?	•
٥	Soil test, other	10	10	۱.
antii	Spraying out pasture including glyphosate	75	75	•
Pre-planting	Lime including cartage and application ²	170	0	•
P	Base fertiliser cost including application	375	0	•
	Cultivation to planting specifications ³	520	520	►
	Pioneer® brand P9978 maize seed @1.30* or 1.35** bags/ha	665	695	Þ
ð	FAR maize seed levy (\$8.00/80,000 kernels @ 1.30* or 1.35** bags/ha)	10	10	•
Planting	LumiGEN® System L-400 seed treatment @ 1.30# or 1.35## bags/ha	170	180)
Ē	Starter fertiliser cost including application ²	295	0	•
	Planting	220	220	►
p	Pre emergence weed control (herbicide + application)	120	120	•
antii	Post emergence weed control (herbicide + application)	105	105	Þ
Post-planting	Sidedress nitrogen cost including application ²	330	0	•
Po	Interest on maize expenditure (7 months @ 6.5%)	135	90	•
	Total growing cost	\$3,200	\$2,025	

Harvest Costs

	Pioneer [®] brand 11C33RR maize specific inoculant ⁴	240	240 270	► ►
Total harvest cost \$1,915 \$1,915 ►	Total harvest cost	\$1,915	\$1,915	۱.

Total Growing & Harvest Costs

\$5,115 \$3,940	\$5,115 \$3,940	\$
-----------------	-----------------	----

*Rounded to the nearest five dollars # Typical fertility ## High fertility



Maize silage drymatter cost

Research has shown that maize can be grown in high fertility dairy farm paddocks, including those with a history of effluent application, without the need for additional fertiliser.

The table below gives indicative drymatter costings for both high and typical fertility maize growing environments. High fertility environments include dairy paddocks coming out of long-term pasture as well as paddocks with a history of effluent application. Typical fertility environments include run-out pasture paddocks and repeat cropping blocks. Very low fertility paddocks including repeat cropping blocks are likely to require additional fertiliser nutrients depending on a number of factors including maize crop yield and winter management system. Always soil test maize paddocks and apply nutrients according to the results.

Drymatter yield per hectare and cost per kg drymatter and per megajoule of metabolisable energy

		Maize silage drymatter costs						
		Typical fertil	ity paddock	High fertility paddock				
	tDM/ha	Maize silage cost per kgDM in the stack (c/kgDM)	Maize silage cost per MJME (c/MJME) ⁵	Maize silage cost per kgDM in the stack (c/kgDM)	Maize silage cost per MJME (c/MJME) ⁵			
	16	32.0	2.96	-	-			
T v	18	28.4	2.63	21.9	2.03			
yiel	20	25.6	2.37	19.7	1.82			
Maize silage yield tDM) in the stack	22	23.3	2.15	17.9	1.66			
ze sil 1) in	24	21.3	1.97	16.4	1.52			
Maiz (†DN	26	19.7	1.82	15.2	1.40			
	28	18.3	1.69	14.1	1.30			
	30	-	-	13.1	1.22			

Assumptions

- 1 Average land rentals have not been included because of a large regional variation. Provision to consider land rental has been included in the My Costs column.
- 2 Fertiliser and lime application rates vary according to soil pH and nutrient status and crop yield targets. Always soil test maize paddocks and seek professional advice to develop a nutrient application plan.
- 3 Cultivation costs will vary depending on soil types, land class and cropping history.
- 4 Costs for Pioneer® brand 11C33RR are based on inoculating a 22 tDM/ha crop.
- 5 Maize silage cost per MJME assumes a maize silage energy content of 10.8 MJME/kgDM.
- 6 Farmers growing maize silage for sale are usually responsible for costs up to and including the sidedress nitrogen application.
- 7 The amount of pasture lost during the maize growing season will vary between paddocks, farms and districts. The value of pasture lost during the maize growing season has not been considered in the calculation of the maize silage drymatter cost.
- 8 The costs and benefits of regrassing have not been included.

Notes

The information in this cost guide is general in nature and is not intended to be representative of actual costs. We do not accept any responsibility or liability (whether as a result of negligence or otherwise) for any loss of any kind that may arise from actions based on the contents of this cost guide or otherwise in connection with the use of this cost guide.

Trait characteristic notes (for table on page 62).

¹ Silage comparative relative maturity (CRM):

Pioneer silage CRM ratings provide a comparison between Pioneer hybrids indicating the relative rates at which hybrids reach harvestable whole plant drymatter. They do **not** represent actual calendar days from planting to harvest.

² Yield for maturity:

Pioneer hybrid trait comparisons should only be made within a range of + or -4 CRM. Analysis of differences in harvest drymatter percentages between hybrids measured in our New Zealand silage research programme show products compared within + or -4 CRM will reach ideal silage harvest maturity (defined as 30% - 38% DM) within about seven days of each other.

³ Adaptability to high population:

A measure of the mix of genetic factors that permit a maize plant to withstand the stresses of high population and still give good standability and high yields.

⁴ Adaptability to low population:

An indicator of a hybrid's ability to compensate (flex) cob size for stand loss from insect damage or poor emergence.

⁵ Stress emergence:

These ratings are based on data collected from local replicated small plot trials planted early and at depth into wet and cold conditions and vigour ratings on New Zealand commercial seed provided by Pioneer's regional seed laboratory. All seed supplied to the market is expected to establish excellent plant stands if planted well and under normal germination conditions. Ratings of 7-9 indicate very good potential to establish normal stands under stressful environmental conditions of cold, wet soils. A 5-6 rating indicates good potential to establish normal stands under moderate stress conditions; and ratings of 1-4 indicate the hybrid has below average potential to establish normal stands under stress and should not be used if severe wet and cold conditions are expected after planting. Hybrids with high ratings are best adapted to early planting but due care to apply best agronomic practices is still required.

⁶ Early growth:

Ratings are taken when two leaf collars are visible.

- ⁷ Plant height:
 - 9 = Tall. 1 = Short.

⁸ Staygreen:

A measure of late season plant health. A lower score means that the plant stover loses colour and dries down more rapidly at maturity.

[°] Whole plant digestibility:

Based on estimated 24 hour in vitro, whole plant digestibility percentage (DM basis) as predicted by Near Infrared Reflectance Spectroscopy (NIRS). A 1 rating point difference reflects 0.50 percent difference in digestibility.

¹⁰ Starch and sugar:

Based on total starch and sugar content of hybrids harvested at silage maturity. Use this score as a relative comparison of the whole plant concentration of readily available energy (primarily grain) among individual hybrids. A 1 rating point difference reflects 0.75 percent difference in starch and sugar.

¹¹ Northern Leaf Blight (NLB) and Eyespot:

In conditions where NLB and Eyespot risks are high, growers should consider planting hybrids with resistance ratings of 6 or higher for these diseases.

¹² Hybrid disease resistance ratings:

8 to 9 = Highly resistant. 6 to 7 = Resistant. 4 to 5 = Moderately resistant. 1 to 3 = Susceptible. - = Insufficient data. Common Rust, Eyespot and NLB ratings are based on overseas data together with New Zealand observations. Scores are based on visual assessment only and not on yield reduction data.

¹³ Cool environments:

In cool environments, including high altitude sites greater than approximately 150 m / 500 ft above sea level, select your yield environment using the descriptions in note 16, then increase established plant populations to the next level i.e. for P9911 in a medium yield environment at high altitude, plant to achieve 115,000 established plants per hectare.

¹⁴ Established plant populations:

These assume good seed establishment conditions. If you are planting very early or into a less than ideal seedbed or where insect pressure may be high (e.g. a shorter than optimum fallow period), seeding rates may need to be increased to compensate for reduced establishment due to field losses.

¹⁵ Plant populations:

The tabulated established populations are recommendations only. Work with your local Pioneer or merchant seed representative to determine the appropriate plant populations for your specific growing environment.

¹⁶ Growing environment definitions:

May include some or all of the following characteristics:

· Challenging yield environments (CYE)

- Typically light, sandy or shallow soils of low fertility, low moisture retention, and predictably low summer rainfall (drought-prone environments).
- High cob or leaf disease pressure.

Medium yield environments (MYE)

- Average fertility soils with predictably adequate summer rainfall and good moisture retention.
- Continuously cropped soils.
- Medium to low cob or leaf disease pressure.

• High yield environments (HYE)

- Typically deeper, highly fertile and well structured soils with good moisture retention.
- Predictably good summer rainfall, shelter from high wind run.
- Good soils straight out of long term pasture.
- Low or no cob or leaf disease pressure.

Publication abbreviations

DM = drymatter bags/ha = bags per hectare kgDM = kilograms of drymatter kgDM/ha = kilograms of drymatter per hectare kgMS = kilograms of milk solids kgMS/cow = kilograms of milksolids per cow \$/ha = dollars per hectare MJME = megajoules of metabolisable energy MJME/kgDM = megajoules of metabolisable energy per kilogram of drymatter t/ha = tonnes per hectare tDM = tonnes of drymatter tDM/cow = tonnes of drymatter per cow tDM/ha = tonnes of drymatter per hectare c/kgDM = cents per kilogram of drymatter



Step by step guide

Choosing the right Pioneer hybrid for your farm

Complete the following four steps to determine the right hybrid for your paddock.



Calculate your days from planting to harvest

Use the chart below to calculate. Line up your planned planting date column with your target harvest date row to find the actual number of days.

•	/	2025						
planting date								
•	/	2026						
harvest date								
•		days						
Days from planting to harvest								
GO TO STEP 2 >								


			Planned planting date 2025														
			Septe	ember				Octobe	r					Nove	mber		
			21	26	1	6	11	16	21	26	31	5	10	15	20	25	30
		5	137	132	127	122	117										
harv	Z	10	142	137	132	127	122	117									
	Februa	15	147	142	137	132	127	122	117								
		20	152	147	142	137	132	127	122	117							
		25	157	152	147	142	137	132	127	122	117						
	March	2	162	157	152	147	142	137	132	127	122	117					
		7	167	162	157	152	147	142	137	132	127	122	117				
		12	172	167	162	157	152	147	142	137	132	127	122	117			
		17	177	172	167	162	157	152	147	142	137	132	127	122	117		
	~	22		177	172	167	162	157	152	147	142	137	132	127	122	117	
		27			177	172	167	162	157	152	147	142	137	132	127	122	117
Target		1				177	172	167	162	157	152	147	142	137	132	127	122
Ta	April	6					177	172	167	162	157	152	147	142	137	132	127
	A	11						177	172	167	162	157	152	147	142	137	132

Notes

It is possible to plant in most areas from late September to mid-December. However, it is important to consider a number of factors that can impact silage yield and quality. These include hybrid pollinating timing relative to potential moisture stress, the planting date needs of the following winter crop or pasture, and the likelihood of autumn frost damage. Please contact your local Pioneer Area Manager or phone 0800 PIONEER (746 633) for advice on hybrid selections for earlier or later plantings.



Use the tables below to choose your region then identify hybrid options by matching your number of days from planting to harvest.



Region 3

South Waikato, King Country, Coastal Taranaki, Rangitikei, Manawatū, Southern Wairarapa & Central Hawke's Bay

Estimated days from

A Contraction of the second se	~
and the second s]
55	
h	

Estimated days from

	Hybrids	planting to harvest		Hybrids	planting to harvest
NEW NEW	P7364	118-133	NEW	P92575	136-154
NEW	P8086	122-136		P9400	140-154
	P8240	124-138	NEW	P9650	144-156
	P8333	125-139		P9911	145-165
	P8666	129-143		P9978	145-161
	P8711	131-145		P0362	147-162

Region 4

Rotorua, Reporoa, Taupo, Central Taranaki, Southern Hawke's Bay, Northern Wairarapa & Horowhenua


A CALLER CONTRACTOR

	Hybrids	Estimated days from planting to harvest	H	ybrids	Estimated days from planting to harvest
NEW	P7179	120-135		8711	139-154
NEW NEW NEW	P7364	127-141	A PS	2575	145-163
NEW	P8086	130-145		9400	149-163
	P8240	132-147		9650	150-165
	P8333	133-148	P	9911	153-173
	P8666	137-152	P	9978	153-169



Notes

Hybrid maturity is based on heat unit accumulation through the season. Hybrids will therefore be quicker to harvest in warmer regions or warmer seasons. For example, a hybrid planted in coastal Nelson will be ready for harvesting much earlier than the same hybrid planted on the same day in mid Canterbury due to the faster accumulation of heat units in the Nelson environment. This has important implications for hybrid selection. The table below is a guide as to the estimated days from planting to harvest for Pioneer® brand hybrids in the regions for which they are recommended for silage. This information is generated from silage hybrid trials carried out in these regions over several seasons.



Region 7

South Canterbury & West Coast

Hybrids

P7179

P7364

P7647

P8086

Estimated days from

planting to harvest

139-153

146-160

148-162

152-166

And	٨

Hybrids

P8666

P8711

Estimated days from planting to harvest P8240 152-167 P8333 153-168

157-172

159-174

Region 8

P8086

North Otago & Southland

		E.
Hybrids	Estimated days from planting to harvest	
P7179	145-160	
P7364	151-166	
P7647	154-168	

156-170



CALCULATORS & GUIDES



		Yield and maturity					Plant traits						Silage quality traits Hybrid disease ratings ¹²			Recommended established plant populations (000s/ ha) ^{13, 14, 15,16}					
	Hybrid	Silage CRM ¹	Grain yield for maturity ²	Silage yield for maturity ²	Adaptability to high population ³	Adaptability to low population ⁴ (ear flex)	Drought tolerance	Stalk strength	Root strength	Stress emergence ⁵	Early growth⁵	Plant height ⁷	Staygreen ^s	Whole plant digestibility 9	Starch and sugar ¹⁰	Northern Leaf Blight ¹¹	Common Rust	Eyespot ^{il}	Challenging yield environments	Medium yield environments	High yield environments
NEW	P7179	71	9	9	9	5	6	7	6	6	7	7	7	8	9	5	6	7	110	120	130
NEW	P7364	73	9	9	9	6	8	7	7	6	7	6	8	9	9	5	4	5	110	120	130
NEW	P7647	76	7	8	6	7	6	7	5	6	7	7	7	9	9	5	6	6	105	110	115
NEW	P8086	80	9	9	7	7	7	6	7	6	6	6	7	7	8	6	7	7	105	115	125
	P8240	82	9	9	7	9	7	6	7	6	6	8	8	7	7	5	5	7	100	110	120
	P8333	83	9	9	5	9	7	6	5	6	8	7	7	7	7	5	6	7	100	110	115
	P8666	86	8	8	5	9	8	6	5	6	8	7	6	6	6	5	6	7	100	110	115
_	P87 11	87	9	8	8	9	8	7	7	6	7	8	8	9	9	7	7	6	100	110	115
NEW	P92575	92	9	9	9	9	8	7	7	7	7	6	9	9	9	8	7	6	95	110	120
	P9400	94	7	7	6	9	7	7	7	4	7	8	6	8	8	7	6	6	100	108	115
NEW	P9650	96	9	9	9	7	7	7	6	6	6	6	7	9	9	7	7	7	95	110	120
	P99 11	99	7	9	7	9	9	5	5	7	6	8	9	6	6	5	5	6	100	108	115
	P9978	99	9	9	9	7	7	6	6	5	6	7	7	9	9	7	7	6	95	110	120
	P0362	103	8	8	9	6	7	6	7	6	6	7	8	7	7	6	7	7	95	105	115
	P0640	106	9	9	7	9	7	6	6	6	8	8	7	7	7	7	7	6	95	105	110
NEW	P0710	107	9	9	9	7	9	7	7	7	8	6	8	8	8	7	7	6	90	100	115
	P0891	107	7	7	9	6	7	8	6	4	6	7	7	7	7	6	6	6	95	105	110
	P0900	109	9	9	9	9	9	7	7	7	7	7	8	7	6	7	7	6	85	95	115
	P0937	109	9	9	9	7	8	6	7	7	8	6	7	8	7	6	6	6	90	100	115
	P1315	110	7	9	5	9	7	7	6	6	7	9	8	5	5	7	7	6	80	90	100
NEW	P1185	111	9	9	9	7	7	8	7	7	7	6	9	8	7	8	8	6	90	100	110
	P1636	112	7	9	8	7	7	6	8	6	7	8	7	6	6	5	6	-	95	105	110
	P1477W	114	9	9	9	6	7	8	8	5	6	7	8	7	7	7	6	7	90	105	110
	P1837	115	7	9	5	9	8	6	6	6	7	9	9	5	5	7	7	7	NR	80	90

Ratings 9 = Outstanding 1 = Poor CRM = Comparative Relative Maturity

– = Insufficient data available NR = not recommended **NEW** = New hybrid For trait notes see page 58







Choose key traits that are important to you

Use trait table on page 62. Enter bybrid trait ratir (right) f the the

the hybrid trait ratings (right) for the hybrids you selected in Step 2.	Hybrid 1	Hybrid 2	Hybrid 3
	•	•	•
	_		_
Silage yield rating ²	•	•	•
Grain yield rating ²	•	•	•
Drought tolerance rating	•	•	•
Staygreen rating ⁸	•	•	•
Early growth rating ⁶	•	•	•
Leaf disease rating ¹¹	•	•	•
Whole plant digestibility ⁹	•	•	•
Starch and sugar ¹⁰	>	•	•
	▶	•	•



Bag calculator

Obtain planting populations from page 62. Determine the number of bags required for each paddock on your farm by completing the table below.

Paddock name

Hectares B X X X * 80 * 80 * 80 Bags required C = = =	Planting population (000's)	Α			
÷ 80 ÷ 80 ÷ 80	Hectares	В	X	X	X
Bags required c = = = =			÷ 80	÷ 80	÷ 80
	Bags required	С	=	=	=

Notes:

•••••	 	 	 	 	 	 	••••••
•••••	 	 	 	 	 •••••	 •••••	•••••••••••
••••••	 	 	 	 	 	 	
•••••	 	 	 	 	 	 	
•••••	 	 	 	 	 	 	••••••
••••••	 	 	 	 	 	 	••••••
••••••	 	 	 	 	 	 	••••••



MATTE KIRK

AREA MANAGER HAMILTON, TAUPIRI & MORRINSVILLE WEST M 027 222 2403 mkirk@genetic.co.nz

SHAUN RUDDELL

AREA MANAGER DARGAVILLE & FAR NORTH M 027 507 4881 sruddell@genetic.co.nz

WARREN COULSON

AREA MANAGER HAURAKI PLAINS, COROMANDEL & MORRINSVILLE EAST M 027 838 7869 wcoulson@genetic.co.nz

GIL DALLAS

AREA MANAGER SOUTH AUCKLAND & NORTH WAIKATO M 027 275 2147 gdallas@genetic.co.nz

COREY THORN

AREA MANAGER WHANGAREI & WELLSFORD M 027 886 0221 cthorn@genetic.co.nz

WADE BELL

FARM SYSTEMS MANAGER NORTHLAND, WAIKATO & BAY OF PLENTY M 027 702 7049 wbell@genetic.co.nz

BRENT BISHOP

REGIONAL MANAGER NORTH WAIKATO & NORTHLAND M 027 554 3315 bbishop@genetic.co.nz



talk to us 0800 746 633



more online at pioneer.nz/contact-us



) facebook.com/pioneerbrandproducts

youtube.com/pioneerbrandproductsnz

Genetic Technologies Limited, Gisborne Office

328 Lytton Road, PO Box 214, Gisborne 4040. Phone: 06 869 0660

Pioneer® brand products are provided subject to the terms and conditions of purchasing, which are part of the labelling and purchase documents *^{10,434} Trademarks and service marks of Dow AgroSciences, DuPont or Pioneer, and their affiliated companies or their respective owners. *²⁰²⁵ Corteva. *²⁰²⁵, Genetic Technologies Limited. All Rights Reserved. The information in this publication is general in nature only. Although the information in this publication is believed to be accurate, no liability (whether as a result of negligence or otherwise) is accepted for any loss of any kind that may arise from actions based on the contents of this publication. *²⁰²⁵, Genetic Technologies Limited. No part of this publication can be reproduced without prior written consent from Genetic Technologies Limited. The form results achieved by testimonial formers are illustrative only of the potential for gains when using Pioneer® brand products. All testimonial figures have been provided and approved by each testimonial farmer. CELEBRATING 50 YEARS IN NEW ZEALAND 1975 - 2025





BEN GORDON

AREA MANAGER SOUTH WAIKATO & CENTRAL PLATEAU M 027 422 7604 bgordon@genetic.co.nz

KAJIL SINGH-SANDHU

AREA MANAGER EASTERN BAY OF PLENTY M 027 220 3848 ksinghsandhu@genetic.co.nz

SOPHIE RIDER

AREA MANAGER KING COUNTRY M 027 214 9084 srider@genetic.co.nz

LOGAN SCOTT AREA MANAGER

TE AWAMUTU EAST & SOUTH WAIKATO M 027 471 0116 Iscott@genetic.co.nz

CRAIG MAXWELL

REGIONAL MANAGER CENTRAL WAIKATO & BAY OF PLENTY M 027 224 0917 cmaxwell@genetic.co.nz

MATT TOWERS

AREA MANAGER TE AWAMUTU WEST M 027 255 3048 mtowers@genetic.co.nz

ROBIN BILLETT

AREA MANAGER EASTERN BAY OF PLENTY M 027 273 0497 rbillett@genetic.co.nz

GRANT DOUGLAS

AREA MANAGER MATAMATA & MORRINSVILLE SOUTH M 027 554 3316 gdouglas@genetic.co.nz

WADE BELL

FARM SYSTEMS SPECIALIST NORTHLAND, WAIKATO & BAY OF PLENTY M 027 702 7049 wbell@genetic.co.nz



talk to us 0800 746 633




more online at pioneer.nz/contact-us



) facebook.com/pioneerbrandproducts

youtube.com/pioneerbrandproductsnz



328 Lytton Road, PO Box 214, Gisborne 4040. Phone: 06 869 0660

Pioneer® brand products are provided subject to the terms and conditions of purchasing, which are part of the labelling and purchase documents. ® ™™ Trademarks and service marks of Daw AgroSciences, DuPont or Pioneer, and their affiliated companies or their respective owners. ® 2025 Corteva. ® 2025, Genetic Technologies Limited. All Rights Reserved. The information in this publication is general in nature only. Although the information in this publication is believed. All Rights Reserved. The information in this publication engligence or otherwise) is accepted for any loss of any kind that may arise from actions based on the contents of this publication. ® 2025, Genetic Technologies Limited. No part of this publication can be reproduced without prior written consent from Genetic Technologies Limited. The farm results achieved by testimonial farmers are illustrative only of the potential for gains when using Pioneer® brand products. All testimonial figures have been provided and approved by each testimonial farmer.







MATT DALLEY

FARMS SYSTEMS SPECIALIST LOWER NORTH ISLAND, TARANAKI & SOUTH ISLAND M 027 508 1719 mdalley@genetic.co.nz

MARK BURKE REGIONAL MANAGER LOWER NORTH ISLAND M 027 451 3987 mburke@genetic.co.nz ALAN BUNNING

SIMON BEGLEY

AREA MANAGER

& EAST COAST

M 027 590 8072

NORTH HAWKE'S BAY

sbegley@genetic.co.nz

AREA MANAGER NORTH TARANAKI M 027 206 0147 abunning@genetic.co.nz KIM SHARPE AREA MANAGER SOUTH TARANAKI M 027 528 0012 ksharpe@genetic.co.nz

CHARLOTTE WILSHER AREA MANAGER HAWKE'S BAY, TARARUA & WAIRARAPA M 027 839 1578 cwilsher@genetic.co.nz DAVID MCDONALD REGIONAL MANAGER TARANAKI, RANGITIKEI & SOUTH ISLAND M 027 276 5439 dmcdonald@genetic.co.nz

RICHARD TEMPLETON AREA MANAGER SOUTHERN MANAWATU & HOROWHENUA M 027 239 0279 rtempleton@genetic.co.nz



talk to us 0800 746 633



more online at pioneer.nz/contact-us youtube.c

) facebook.com/pioneerbrandproducts

youtube.com/pioneerbrandproductsnz

Genetic Technologies Limited, Gisborne Office

328 Lytton Road, PO Box 214, Gisborne 4040. Phone: 06 869 0660

Pioneer® brand products are provided subject to the terms and conditions of purchasing, which are part of the labelling and purchase documents. [®] ML^{MM} Trademarks and service marks of Dow AgroSciences, DUPont or Pioneer, and their affiliated companies or their respective owners. [®]2025 Corteva. [®]2025, Genetic Technologies Limited. All Rights Reserved. The information in this publication is general in nature only. Although the information in this publication is believed to be accurate, no liability (whether as a result of negligence or otherwise) is accepted for any loss of any kind that may arise from actions based on the contents of this publication. [®]2025, Genetic Technologies Limited. No part of this publication can be reproduced without prior written consent from Genetic Technologies Limited. The form results achieved by testimonial farmers are illustrative only of the potential for gains when using Pioneer® brand products. All testimonial figures have been provided and approved by each testimonial farmer.







MATT DALLEY

FARMS SYSTEMS SPECIALIST SOUTH ISLAND, TARANAKI & LOWER NORTH ISLAND M 027 508 1719 mdalley@genetic.co.nz

PAIGE FARRELL AREA MANAGER NORTH CANTERBURY, **TASMAN & MARLBOROUGH** M 027 251 1316 pfarrell@genetic.co.nz

talk to us

0800 746 633

DARYL MOORE

AREA MANAGER **SOUTH CANTERBURY &** NORTH OTAGO M 027 767 1119 dmoore@genetic.co.nz

DUNCAN GILLANDERS

AREA MANAGER **MID CANTERBURY &** WEST COAST M 027 555 9016 dgillanders@genetic.co.nz

DARIN DICK

RACHEL BELL

AREA MANAGER

M 027 839 7657

rbell@genetic.co.nz

CANTERBURY

AREA MANAGER **OTAGO & SOUTHLAND** M 027 546 0005 ddick@genetic.co.nz

DAVID MCDONALD

REGIONAL MANAGER SOUTH ISLAND, **TARANAKI & RANGITIKEI** M 027 276 5439 dmcdonald@genetic.co.nz

facebook.com/pioneerbrandproducts

youtube.com/pioneerbrandproductsnz

Genetic Technologies Limited, Gisborne Office

328 Lytton Road, PO Box 214, Gisborne 4040. Phone: 06 869 0660

more online at

pioneer.nz/contact-us

Pioneer® brand products are provided subject to the terms and conditions of purchasing, which are part of the labelling and purchase documents. ^{e, TM, SM} Trademarks and service marks of Dow AgroSciences, DuPont or Pioneer, and their affiliated companies or their respective owners. "2025 Corteva. "2025, Genetic Technologies Limited. All Rights Reserved. The information in this publication is general in nature only. Although the information in this publication is believed to be accurate, no liability (whether as a result of negligence or otherwise) is accepted for any loss of any kind that may arise from actions based on the contents of this publication. *2025, Genetic Technologies Limited. No part of this publication can be reproduced without prior written consent from Genetic Technologies Limited. The farm results achieved by testimonial farmers are illustrative only of the potential for gains when using Pioneer® brand products. All testimonial figures have been provided and approved by each testimonial farmer.



